arXiv:hep-th/0308146v3 17 Sep 2003

N=1 Supersymmetric Boundary Bootstrap

Gábor Zsolt Tóth

21st August 2003

Institute for Theoretical Physics,
Eötvös University, Budapest, Hungary
H-1117 Budapest, Pázmány Péter Sétány 1/A

`tgabor@afavant.elte.hu`

Abstract

We investigate the boundary bootstrap programme for finding exact reflection matrices of integrable boundary quantum field theories with N=1 boundary supersymmetry. The bulk S-matrix and the reflection matrix are assumed to take the form $\hat{S}=\tilde{S}S$, $\hat{R}=\tilde{R}R$, where $\tilde{S}$ and $\tilde{R}$ are the S-matrix and reflection matrix of some integrable non-supersymmetric boundary theory that is assumed to be known, and S and R describe the mixing of supersymmetric indices. Under the assumption that the bulk particles transform in the kink and boson/fermion representations and the ground state is a singlet we present rules by which the supersymmetry representations and reflection factors for excited boundary bound states can be determined. We apply these rules to the boundary sine-Gordon model, to the boundary $a_2^{(1)}$ and $a_4^{(1)}$ affine Toda field theories, to the boundary sinh-Gordon model and to the free particle.

PACS codes: 11.10.Kk, 11.55.Ds, 11.30.Pb
Keywords: supersymmetry, integrable quantum field theory, boundary scattering, bound state, bootstrap, sine-Gordon model, sinh-Gordon model, affine Toda theory

1 Introduction

Integrable quantum field theories (QFTs) in two dimensions have the remarkable property that their S-matrix factorizes and the bootstrap programme becomes manageable. The particle spectrum and S-matrix of several integrable QFTs have been found by applying the axioms of general S-matrix theory and factorized scattering theory. Studying supersymmetric field theories it is an interesting problem to determine the S-matrix and particle spectrum of integrable supersymmetrized QFTs.

A formalism for constructing $N = 1$ supersymmetric factorizable scattering theory is developed in [3, 1, 17, 6, 5]. The particles of the non-supersymmetric theory are assumed to become supermultiplets $|\,\xi_i(\theta)\rangle \otimes |\,A_i(\theta)\rangle$, where $|\,\xi_i(\theta)\rangle$ are the one-particle states of the non-supersymmetric theory, and it is also assumed that all the particles of the supersymmetric theory are obtained in this way. The index i specifies (not necessarily all) Poincare-invariant conserved quantum numbers of the non-supersymmetric theory and A_i specifies the supersymmetric quantum numbers. Those conserved quantum numbers which are specified by i must not change their value under charge conjugation, and supersymmetry (except for boosts) acts trivially on the $|\,\xi_i(\theta)\rangle$ part of the states. As the simplest Ansatz the product form $\hat{S}^{\xi'_j A'_j;\xi'_i A'_i}_{\xi_i A_i;\xi_j A_j}(\theta_1-\theta_2) = \tilde{S}^{\xi'_j\xi'_i}_{\xi_i\xi_j}(\theta_1-\theta_2) S^{A'_j A'_i}_{A_i A_j}(\theta_1-\theta_2)$ was proposed for the two-particle S-matrix $\hat{S}$, where $\tilde{S}^{\xi'_j\xi'_i}_{\xi_i\xi_j}(\theta)$ is the two-particle S-matrix of the non-supersymmetric theory subject to supersymmetrization, and the supersymmetric factor $S^{A'_j A'_i}_{A_i A_j}(\theta)$ describes the mixing of supersymmetric indices (see also the first paragraph of section 2 for conventions). The requirement that the particle spectrum can be obtained as described above is consistent with the Ansatz for the S-matrix if "minimal" supersymmetric S-matrix factors having no poles and overall zeroes on the imaginary axis in the physical strip are used. We restrict ourselves to such S-matrix factors (see section 5.2 for an exception). As the notation indicates, it is required that the SUSY representations associated to the particles depend on conserved Poincare- and charge conjugation invariant quantum numbers only, what guarantees that the complete S-matrix (including amplitudes of multi-particle scatterings) of the supersymmetric theory also takes a product form. The complete two-particle S-matrix $\hat{S}$ and so the supersymmetric factor S commute with the action of supersymmetry. The Yang-Baxter, fusing, and bootstrap equations, the crossing equation, the unitarity and analyticity conditions are satisfied by $\hat{S}$ and take a factorized form. Since $\tilde{S}$ satisfies them, the supersymmetric factor S must also satisfy them separately. The fusing angles occurring in the fusing and bootstrap equations for S are determined by the non-supersymmetric theory, and since the bound state poles are already present in $\tilde{S}$, the supersymmetric factor S satisfies the bootstrap equations passively, i.e. S does not have poles at the fusing rapidities.

An essential step in this construction is the choice of SUSY representations in which the particles transform. Having made this choice the supersymmetric factor S of the two-particle S-matrix is obtained by solving the supersymmetry condition, the Yang-Baxter and crossing equations, the unitarity, analyticity and minimality conditions. These equations depend on the conserved quantum numbers, so they impose restrictions on the choice of representations. Further highly nontrivial restrictions come from the bootstrap and especially from the fusing equations for S, which contain the fusing data of the non-supersymmetric model. Let $a_i + b_j \to c_k$ be a fusion process of the non-supersymmetric theory with fusing rapidity $iu^{c_k}_{a_i b_j}$, and assume that the representations in which a_i, b_j, and c_k transform are D_i, D_j and D_k; in this case the fusing equation is equivalent to the statement that the supersymmetric two-particle S-matrix block at rapidity $iu_{a_ib_j}^{c_k}$ is an equivariant projection from the product of D_i and D_j on D_k. This can be very restrictive for the value of $u_{a_ib_j}^{c_k}$, as in the most common case, for example, when D_i, D_j, D_k are boson/fermion representations [1, 3]. We remark that the fusing equation can be forced to hold in certain cases by factoring out unwanted states from the Hilbert-space.

The previous considerations indicate that it is nontrivial that the construction above can be applied to supersymmetrize a given factorized scattering theory. The knowledge of the Lagrangian density or other data of the (supersymmetric) theory often suggests preferred SUSY representations for the particle multiplets. If the possible representations in which the particles can transform are fixed, then by solving the axioms above one can derive necessary and sufficient conditions that have to be satisfied by the particle spectrum and fusing rules of a non-supersymmetric theory which is supersymmetrized using these representations. These conditions have been obtained in the case when the possible representations are the kink and the boson/fermion representation, and several factorizable scattering theories —the $a_{n-1}^{(1)}$, $d_n^{(1)}$, $(c_n^{(1)}, d_{n+1}^{(2)})$ and $(b_n^{(1)}, a_{2n-1}^{(2)})$ affine Toda theories, the $SU(2)$ principal chiral model, the sine-Gordon model, the supersymmetric $O(2n)$ sigma model [1], and the multicomponent Yang-Lee (or FKM) minimal models [4]— have been found to satisfy these conditions, i.e. these theories have been supersymmetrized by applying the construction described above. The Lagrangian field theories underlying these supersymmetric scattering theories are not all known (if they exist).

It is a natural step after the study of bulk factorized scattering theories to study them in the presence of a boundary. If the boundary conditions and interactions preserve integrability, then the bootstrap programme remains manageable. A set of boundary bound states are added to the bulk spectrum, and scattering processes involving boundary states are described by reflections amplitudes which can be written as products of one-particle reflection amplitudes and two-particle S-matrices. The boundary versions of the unitarity condition, crossing equation, Yang-Baxter, fusing, and bootstrap equations can be introduced [19]. The Ansatz above for constructing supersymmetric scattering theories can also be extended to the situation when an integrable boundary is present. A nontrivial problem of this extension that we discuss in this paper is the definition of the concept of supersymmetry in the presence of boundary. Results in the classical Lagrangian field theory framework [7] indicate clearly that such a concept is possible. Our construction is based on the results in [15, 16, 2, 4, 7, 8] and it can be regarded as an application of the construction described in [15, 16] for quantum group symmetry.

Let us assume that a factorizable scattering theory with boundary is given and the supersymmetric version of the bulk part of this theory is also constructed using the Ansatz above, the bulk particles transforming either in the kink or in the boson/fermion representation. We study in this paper the problem of completing the supersymmetrization in this case using the boundary version of the Ansatz above. The first step is to choose a representation of the boundary supersymmetry algebra for the ground state. Next the minimal supersymmetric factors of the ground state one-particle reflection amplitudes are to be determined using the boundary Yang-Baxter, unitarity and crossing equations and the supersymmetry condition for these factors. Finally the boundary bootstrap and fusing equations for supersymmetric factors can be used to obtain the representations in which the excited boundary bound states transform together with the supersymmetric factors of the one-particle reflection amplitudes on these states. We remark that if there were preferred representations for the excited boundary states, then the boundary fusing equation would constrain the boundary fusing angles. The first and especially the second steps have been considered in the literature [14, 10, 4, 9, 8, 2], whereas there are fewer results concerning the last step [2]. There are few known solutions to the boundary bootstrap programme for non-supersymmetric theories either.

For the ground state we take the singlet representations with RSOS label $\frac{1}{2}$, this being the simplest case and also because we expect [2, 14] that other cases can be obtained by boundary bootstrap from this one. The general minimal supersymmetric one-particle ground state reflection factors have been determined for this case in [14, 10, 9] but without imposing the supersymmetry condition. This condition is imposed on the kink reflection amplitude in [2]. In the present paper we consider the boson/fermion ground state reflection factors in some detail. We derive these reflection factors by imposing the supersymmetry condition and solving the boundary Yang-Baxter equation afterwards.

As the main result of the paper we determine the representations and supersymmetric one-particle reflection factors for excited boundary bound states using the bootstrap and fusing equations, for arbitrary boundary fusing rules. This is a nontrivial problem because the one-particle reflection amplitudes and so the fusing tensors can be degenerate at particular rapidities. Although these rapidities are special, they usually play an essential role in boundary theories. We also present the conditions on the fusing angles resulting from the requirement of minimality. This requirement is nontrivial even if the ground state reflection factors are minimal, and restricts the boundary fusing angles. Our work is motivated by the results of [1] for bulk supersymmetric bootstrap, and by [2], where the scattering theory of the boundary sine-Gordon model (BSG) is supersymmetrized. In the BSG theory the whole boundary spectrum can be generated by kinks, and in this case it is not difficult to determine the representations and supersymmetric reflection factors for the boundary states. However, there are other ways to generate the boundary states, and it is nontrivial —though plausible— that the same representations and reflection factors are obtained in each way. The correctness of this proposition for the BSG model was conjectured and partially verified in [2]. Using our results we can complete the verification and confirm that the proposition is correct.

Only few integrable boundary scattering theories are known explicitely. Beyond the BSG model non-trivial scattering theories for boundary $a_2^{(1)}$ and $a_4^{(1)}$ affine Toda field theories [13], for the free boson on the half line, and for the boundary sinh-Gordon model [12] are known. The bulk part of these theories can be supersymmetrized in the framework described above, so we apply our results to them.

The layout of the paper is as follows. In section 2 we review the formalism for supersymmetric factorized scattering theory in the bulk as described in [3], [1] and [17]. This review is not a mere copy of results as we found the thorough understanding of this formalism necessary for studying the boundary case. We hope that we make certain points clearer, and we also supply certain details (concerning representation theory for instance) that cannot be found in [3, 1, 17]. (The first part of section 4.4 is also relevant for the bulk formalism.) In section 3 we discuss the boundary supersymmetry algebra and its action on the ground state and multi-particle states, the boundary version of the Ansatz for supersymmetric scattering theory, and the supersymmetric one-particle ground state reflection factors for kinks and boson/fermion doublets. Section 4 contains a description of the boundary supersymmetric bootstrap structure and fusing rules under the condition that the bulk particles transform in the kink and boson/fermion representations and the ground state is a singlet with RSOS label $\frac{1}{2}$. This section and especially subsection 4.4 contains the main results of the paper also mentioned above. In section 5 we apply the general results of the previous sections to specific models. We present our conclusions in section 6. The Appendix contains the normalization factors for the S-matrix blocks.

2 Bulk supersymmetric factorized scattering and bootstrap

The various particles contained in multi-particle *in, out* or *intermediate* states will be written in their spatial order. The order of the indices of the S-matrices will also be the same as the spatial order of the corresponding particles (see figure 1). On the figures time flows upward. Upper indices of the S-matrix will correspond to *out* states: $|\,c\rangle(\theta_1)d(\theta_2)_{in} = S^{ab}_{cd}(\theta_1-\theta_2)|\,a\rangle(\theta_2)b(\theta_1)_{out}$. If the S-matrix is written in matrix form (i.e. as a table of entries), then the upper indices specify the rows and the lower indices specify the columns. These conventions agree with [3]. The "physical strip" for the rapidity argument θ of the S-matrix is $0 < Im(\theta) < \pi$.

Throughout the paper we mainly deal with the supersymmetric factors of the S-matrix and with the supersymmetric part $|\,A_i(\theta)\rangle$ of the states. We shall usually refer to $|\,A_i(\theta)\rangle$ itself as a state or supersymmetric state.

2.1 Bulk supersymmetry charges

The generators of the supersymmetry algebra are denoted by N, Q, $\bar{Q}$, Γ. Q and $\bar{Q}$ are called supercharges, Γ is the fermionic parity operator and N is the boost generator. The generators Q, $\bar{Q}$, Γ satisfy the relations

$$\Gamma^2 = 1, \qquad \{\Gamma, Q\} = \{\Gamma, \bar{Q}\} = 0. \tag{1}$$



Figure 1: Two-particle S-matrix

The translation generators H and P are related to the supercharges:

$$Q^2 = H + P, \qquad \bar{Q}^2 = H - P. \tag{2}$$

It follows that the SUSY central charge $\hat{Z} = \frac{1}{2}\{Q, \bar{Q}\}$ commutes with Q, $\bar{Q}$, Γ, H and P. There are also the relations

$$[N, H+P] = H+P, \quad [N, H-P] = -(H-P), \quad [N, Q] = \frac{1}{2}Q, \quad [N, \bar{Q}] = -\frac{1}{2}\bar{Q}. \tag{3}$$

The coproduct Δ used to define the action of the supersymmetry algebra on multi-particle states is given by

$$\Delta(Q) = Q \otimes I + \Gamma \otimes Q, \quad \Delta(\bar{Q}) = \bar{Q} \otimes I + \Gamma \otimes \bar{Q}, \tag{4}$$

$$\Delta(\Gamma) = \Gamma \otimes \Gamma, \quad \Delta(Z) = \hat{Z} \otimes I + I \otimes \hat{Z},$$

$$\Delta(H) = H \otimes I + I \otimes H, \quad \Delta(P) = P \otimes I + I \otimes P, \quad \Delta(N) = N \otimes I + I \otimes N.$$

On a one-particle supersymmetric multiplet $\mid A_i\rangle$

$$Q|\, A_i(\theta)\rangle = \sqrt{m_i} e^{\theta/2} q|\, A_i(\theta)\rangle, \quad \bar{Q}|\, A_i(\theta)\rangle = \sqrt{m_i} e^{-\theta/2} \bar{q}|\, A_i(\theta)\rangle,$$

where m_i is the mass of the multiplet and q and $\bar{q}$ are θ-independent matrices which act on the states of the supermultiplet and satisfy

$$q^2 = 1, \qquad \bar{q}^2 = 1, \qquad \{q, \bar{q}\} = 2Z,$$

where $Z = \frac{1}{m_i}\hat{Z}$ on the multiplet. The action of Γ is independent of θ and

$$\{\Gamma, q\} = \{\Gamma, \bar{q}\} = 0.$$

2.2 Particle and kink representations

Two representations of the supersymmetry algebra are used to construct supermultiplets: the boson-fermion and the kink representation. The boson-fermion representation will often be referred to as the particle representation. It contains a boson $|\,\phi\rangle$ and a fermion $|\,\psi\rangle$. In the basis $\{\phi,\psi\}$

$$q=\begin{pmatrix} 0 & \epsilon \\ \epsilon^* & 0 \end{pmatrix},\quad \bar{q}=\begin{pmatrix} 0 & \epsilon^* \\ \epsilon & 0 \end{pmatrix},\quad \Gamma=\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \tag{5}$$

where $\epsilon=\exp(i\pi/4)$. The central charge is zero. Charge conjugation C acts as the identity: $\phi\leftrightarrow\phi$, $\psi\leftrightarrow\psi$. We denote this representation by P. We will often refer to bosons and fermions simply as particles.

Another representation is obtained if we multiply Γ in (5) by -1. We call it pseudoparticle representation and denote it by $\bar{P}$. The following equations describe the decomposition of two-particle states:

$$P\otimes P=P\oplus\bar{P},\quad \bar{P}\otimes\bar{P}=P\oplus\bar{P},\quad P\otimes\bar{P}=P\oplus\bar{P}. \tag{6}$$

In (6) the third equation means for example that a two-particle state containing a particle and a pseudoparticle transforms in the sum of a particle and a pseudoparticle representation (of appropriate mass).

The kink representation K contains four kinks, interpolating between three vacua labeled by the RSOS labels $0,\frac{1}{2},1$. In the basis $\{K_{0\frac{1}{2}},K_{1\frac{1}{2}},K_{\frac{1}{2}0},K_{\frac{1}{2}1}\}$

$$q=\begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},\quad \bar{q}=\begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},\quad \Gamma=\begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \tag{7}$$

The central charge is

$$Z=\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$

There is also another representation $\bar{K}$ called pseudo-kink representation, which is obtained by multiplying q and $\bar{q}$ by -1 in (7), or by interchanging the RSOS labels $0\leftrightarrow 1$. Multi-kink states have to respect an adjacency condition: in the physical $|\,...K_{ab}(\theta_1)K_{cd}(\theta_2)...\rangle$ state $b=c$ must hold. This condition gives kinks a character rather different from that of particles. (The condition also implies that $(\Gamma\otimes I)|\,K_{\frac{1}{2}1}(\theta_1)K_{1\frac{1}{2}}(\theta_2)\rangle=0$, for example.) Charge conjugation acts as follows: $K_{0\frac{1}{2}}\leftrightarrow K_{\frac{1}{2}0}$, $K_{1\frac{1}{2}}\leftrightarrow K_{\frac{1}{2}1}$. Multi-kink states containing even number of kinks can be arranged in two sectors: the first sector contains the states which have left and right RSOS index $\frac{1}{2}$, the second sector contains the states which have left and right RSOS indices 0 or 1. These two sectors will be called $\frac{1}{2}$ and 01 sector. For two-kink states we have

$$K \otimes K = [P]_{\frac{1}{2}} \oplus [P \oplus \bar{P}]_{01}. \tag{8}$$

The subscripts refer to the sectors in which the subspaces lie.

The $\otimes$ symbol in (8) and further on denotes a tensor product with the modification that the adjacency conditions (if there are any) are also understood to be imposed. Thus $\otimes$ will sometimes stand for a non-free tensor product, but this will not be denoted explicitly.

The following combinations of two-kink states span the invariant subspaces (see also [17, 10]):

$$|\,\phi_1\rangle = |\,K_{\frac{1}{2}0}K_{0\frac{1}{2}}\rangle + |\,K_{\frac{1}{2}1}K_{1\frac{1}{2}}\rangle, \qquad |\,\psi_1\rangle = |\,K_{\frac{1}{2}0}K_{0\frac{1}{2}}\rangle - |\,K_{\frac{1}{2}1}K_{1\frac{1}{2}}\rangle; \tag{9}$$

$$|\,\phi_2\rangle = |\,K_{0\frac{1}{2}}K_{\frac{1}{2}0}\rangle + |\,K_{1\frac{1}{2}}K_{\frac{1}{2}1}\rangle, \qquad |\,\psi_2\rangle = |\,K_{1\frac{1}{2}}K_{\frac{1}{2}0}\rangle - |\,K_{0\frac{1}{2}}K_{\frac{1}{2}1}\rangle; \tag{10}$$

$$|\,\bar{\phi}\rangle = |\,K_{0\frac{1}{2}}K_{\frac{1}{2}0}\rangle - |\,K_{1\frac{1}{2}}K_{\frac{1}{2}1}\rangle, \qquad |\,\bar{\psi}\rangle = |\,K_{1\frac{1}{2}}K_{\frac{1}{2}0}\rangle + |\,K_{0\frac{1}{2}}K_{\frac{1}{2}1}\rangle. \tag{11}$$

The rapidities do not play essential role in these formulae, so they are suppressed. The two states in (11) span the pseudoparticle representation. The value of Γ on $|\,\bar{\phi}\rangle$ is -1. Although the decomposition (8) is possible, the products of elements of $[P]_{\frac{1}{2}}$, $[P]_{01}$ and $[\bar{P}]_{01}$ satisfy certain relations because of the kink adjacency conditions. For example, $|\,\bar{\phi}\bar{\phi}\rangle = |\,\phi_2\phi_2\rangle$.

The eight particle-kink states $|\,p(\theta_1)k(\theta_2)\rangle$, where p stands for a boson or fermion and k stands for a kink, transform in the direct sum of a kink and a pseudo-kink representation if and only if

$$\theta_1 - \theta_2 = i(\pi - u) \quad \text{and} \quad m = 2M\cos(u) = 2M\sin(\pi/2 - u), \tag{12}$$

where m is the mass of the particle and M is the mass of the kink. This is precisely the condition that the total mass of the particle-kink state is also M. If this condition is not satisfied, then the decomposition of the representation in which the particle-kink states transform does not contain the kink, pseudo-kink, particle or pseudoparticle representations.

2.3 Scattering amplitudes and supersymmetric bootstrap

The general solution of the Yang-Baxter equations that describes the (supersymmetric factor of the) scattering of particle supermultiplets is

$$S_P^{[i,j]}(\theta) = G^{[i,j]}(\theta)\left[\frac{1}{2i}(q_1 - q_2)(\bar{q}_1 - \bar{q}_2) + \alpha F(\theta)[1 - t(\theta, m_i, m_j)q_1 q_2][1 + t(\theta, m_j, m_i)\bar{q}_1\bar{q}_2]\right], \tag{13}$$

where

$$t(\theta,m_i,m_j)=\tanh\left(\frac{\theta+\log(m_i/m_j)}{4}\right),\qquad F(\theta)=\frac{m_i+m_j+2\sqrt{m_im_j}\cosh(\theta/2)}{2i\sinh(\theta)},$$

$q_1=q\otimes I$, $q_2=\Gamma\otimes q$, $\bar{q}_1=\bar{q}\otimes I$, $\bar{q}_2=\Gamma\otimes\bar{q}$. m_i and m_j are the masses of the multiplets, and α is a real constant, which measures the strength of Bose-Fermi mixing (the dependence of α on i,j is not indicated). $S_P^{[i,j]}(\theta)/G^{[i,j]}(\theta)$ can depend on the conserved quantities i,j through m_i, m_j and α only. It also follows from the Yang-Baxter equation that the particles in a theory can be divided into disjoint sets with the property that any two particles in a set have the same nonzero α, and $\alpha=0$ for two particles from different sets. $\alpha=0$ corresponds to trivial scattering. To each particle in a theory we associate a value of α, which is the value that occurs in the scattering of the particle with itself. For simplicity we consider only theories which have only one such set and thus α is the same for any two-particle scattering. The scalar function $G^{[i,j]}(\theta)$ is determined by unitarity and crossing symmetry up to CDD factors. It is important here that i,j are invariant under charge conjugation. It is also required that $S_P^{[i,j]}(\theta)$ should be minimal, what fixes $G^{[i,j]}(\theta)$ completely. The minimal solution can be found in the Appendix (cf. [1, 3]). $G^{[i,j]}(\theta)$ contains the parameters u_i, u_j for which

$$0<Re(u_i),Re(u_j)\leq\pi/2,\qquad m_i=2M\sin(u_i),\quad m_j=2M\sin(u_j),\tag{14}$$

where $M=|1/(2\alpha)|$. Consequently, for each particle there is a corresponding angle u. In this paper we consider only real values of u_i and u_j.

The S-matrix factor that describes the scattering of two kinks of equal mass is

$$S_K^{[i,j]}(\theta)=K^{[i,j]}(\theta)[\cosh(\gamma\theta)-\sinh(\gamma\theta)q_1\bar{q}_1][\cosh(\theta/4)-\sinh(\theta/4)q_1q_2],$$

where $\gamma=\log 2/2\pi i$. The scalar function $K^{[i,j]}(\theta)$ is determined by unitarity and crossing symmetry and the condition that $S_K^{[i,j]}(\theta)$ should be minimal. See the Appendix or [1] for $K^{[i,j]}(\theta)$. $S_K^{[i,j]}(\theta)$ is bijective in the physical strip. The scattering of kinks of different mass is impossible, so all the kinks in a theory have to have the same mass. The kink-particle S-matrix $S_{PK}(\theta)$ will be considered later. The upper indices ${}^{[i,j]}$ will often be suppressed.

The important common feature of these minimal S-matrices, including S_{PK}, is that they have no poles and overall zeroes in the physical strip (although they can be degenerate at particular values of θ). It is required that the scattering of all particles and kinks of the supersymmetrized theory should be described by minimal supersymmetric S-matrix factors, i.e. S_P, S_K and S_{PK} must be the supersymmetric two-particle S-matrix building blocks. It turns out that this requirement is consistent with the bootstrap equation, so the fusing data of the non-supersymmetric theory are not modified essentially by adding the supersymmetric S-matrix factors.



Figure 2: The fusing equation (splitting of S)

The three-particle couplings (or fusing and decay tensors) are also assumed to factorize into a non-supersymmetric and supersymmetric part: $\hat{f}^{\xi_k A_k}_{\xi_i A_i \xi_j A_j} = \tilde{f}^{\xi_k}_{\xi_i \xi_j} f^{A_k}_{A_i A_j}$, $\hat{d}^{\xi_i A_i \xi_j A_j}_{\xi_k A_k} = \tilde{d}^{\xi_i \xi_j}_{\xi_k} d^{A_i A_j}_{A_k}$, and for the supersymmetric factor of the S-matrix

$$S^{A'_j A'_i}_{A_i A_j}(iu^k_{ij}) = \sum_{A_k} f^{A_k}_{A_i A_j} d^{A'_j A'_i}_{A_k} \tag{15}$$

holds, where u^k_{ij} denotes the $\xi_i + \xi_j \to \xi_k$ fusing angle, and

$$|\, A_i(\theta + i(\pi - u^j_{ik}))A_j(\theta - i(\pi - u^i_{jk}))\rangle = \sum_{A_k} f^{A_k}_{A_i A_j} |\, A_k(\theta)\rangle, \tag{16}$$

$$|\, A_k(\theta)\rangle = \sum_{A_i, A_j} d^{A_i A_j}_{A_k} |\, A_i(\theta - i(\pi - u^j_{ik}))A_j(\theta + i(\pi - u^i_{jk}))\rangle, \tag{17}$$

where u^j_{ik}, u^i_{jk} are the other angles at the three-particle vertices. We call these equations fusing equations. The f fusing tensors and the d decay tensors should commute with the action of supersymmetry. See fig. 2 for graphical representation.

Once the supersymmetry representations in which the various particles transform are fixed, these equations can severely constrain the possible fusing angles in a sense that we describe next, beginning with $particle + particle \to particle$ fusion.

In the light of (6), if it is decided that some particles and their bound states transform in the particle representation of the SUSY algebra, then the fusing equation (see figure 2) can be satisfied if and only if $S_P(iu^k_{ij})$ is a projection on the appropriate subspace carrying the particle representation. This is a nontrivial condition on $S_P(iu^k_{ij})$, because $S_P(\theta)$ is bijective (of rank four) in general. The other possibility to assure that only particles are produced in the fusion is to factor out the unwanted states by hand. Note that because of (16) and (17) this implies that certain two-particle *in* and *out* states with complex rapidities also have to be factored out. In the present paper we consider the first, more natural possibility.

$S^{[i,j]}_P(\theta)$ is of rank two if $\theta = iu^k_{ij}$, where $u^k_{ij} \in \{u_i + u_j, \pi - u_i + u_j, u_i + \pi - u_j\}$. Only two of these values can be in the physical strip, and $S^{[i,j]}_P(\theta)$ is nondegenerate at



Figure 3: Bootstrap equation

other values of θ in the physical strip. The image space of $S_P^{[i,j]}(iu_{ij}^k)$ carries the particle representation if and only if $\alpha < 0$, i.e. if $\alpha = -1/(2M)$. We remark that if $\alpha = 1/(2M)$, then the image space carries the pseudoparticle representation. The value of u_k is the following:

$$u_k = u_i + u_j \quad \text{if} \quad u_{ij}^k = u_i + u_j < \pi/2, \tag{18}$$

$$u_k = \pi - (u_i + u_j) \quad \text{if} \quad u_{ij}^k = u_i + u_j \geq \pi/2, \tag{19}$$

$$u_k = u_i - u_j \quad \text{if} \quad u_{ij}^k = \pi - u_i + u_j, \tag{20}$$

$$u_k = u_j - u_i \quad \text{if} \quad u_{ij}^k = u_i + \pi - u_j. \tag{21}$$

The other angles at the three-particle vertex are

$$\begin{aligned} &u_{jk}^i = \pi - u_i, \quad u_{ki}^j = \pi - u_j \quad \text{if} \quad u_{ij}^k = u_i + u_j, \\ &u_{jk}^i = u_i, \quad u_{ki}^j = \pi - u_j \quad \text{if} \quad u_{ij}^k = \pi - u_i - u_j, \\ &u_{jk}^i = \pi - u_i, \quad u_{ki}^j = u_j \quad \text{if} \quad u_{ij}^k = u_i + \pi - u_j. \end{aligned} \tag{22}$$

The fusions corresponding to (20) or (21) are crossed versions of (18), (19). Explicit expressions for the three-particle couplings are written down in [3, 1].

As discussed above, we require the minimality of the supersymmetric two-particle S-matrices. This is yet another nontrivial condition for the possible fusions, because the bootstrap equation determines the scattering of the produced particle on other particles. It is clear that the (two-particle) supersymmetric S-matrix given by the bootstrap equation will be S_P with $\alpha = -1/(2M)$ modulo CDD factors, because a two-particle S-matrix given by the bootstrap equation automatically satisfies the axioms of factorized scattering, i.e. the Yang-Baxter, unitarity and crossing equations, and real analyticity, and we know that the solution to these constraints is S_P modulo CDD factors. By checking the pole structure it can be seen that the (supersymmetric) particle S-matrix given by bootstrap is in fact also S_P, so the minimality condition is consistent with the bootstrap and does not give more constraints on the fusion of (supersymmetric) particles.

We turn to the case of the fusion of two (supersymmetric) kinks of equal mass now. The kink scattering amplitude is always of rank six (bijective) in the physical strip. Consequently, in the light of (8), if one insists that no pseudoparticles should be formed in kink fusion, then one has to factor out the unwanted states from the Hilbert-space by hand. Because of (16) and (17) this implies that certain two-kink *in* and *out* states with complex rapidities also have to be factored out. Even if states of the form (11) with kink rapidities $\theta + iu$ and $\theta - iu$, $2u$ being the fusing angle, are factored out, kink fusion produces two types of particles corresponding to (9) and (10), i.e. to the $\frac{1}{2}$ and 01 sectors. The two types of particles will be referred to as type $\frac{1}{2}$ and type 01 particles.

There are adjacency conditions for particles produced in kink fusion, which follow from the adjacency conditions for kinks: type $\frac{1}{2}$ and type 01 particles cannot be adjacent in a multi-particle state, so they cannot scatter on each other. There are also appropriate adjacency conditions for kinks and particles. Bootstrap gives the result [17, 1] that the two types of particles have the same S-matrix S_P. Consequently, the two types of particles can be identified (which is the same as factoring out certain combinations). If this identification is made, then only the following adjacency condition applies: if in a state $|\, ...K_{ab}p...p...pK_{cd}...\rangle$ (where p stands for a particle and the rapidities are suppressed) there are only particles between K_{ab} and K_{cd}, then either $b = c$, or $|b - c| = 1$.

In this paper we restrict ourselves to theories in which pseudoparticles do not occur (see also section 4.4). Consequently, we have to factor out the pseudoparticles that would arise. We also identify the two types of particles[1].

There is no condition on the rank of the kink-kink scattering amplitude $S^{A'_j A'_i}_{A_i A_j}(iu^k_{ij})$, and so the fusing equations can be satisfied for arbitrary fusing angles. The mass of the resulting state will be

$$m_3 = 2M\cos(u^k_{ij}/2) = 2M\sin(\pi/2 - u^k_{ij}/2),$$

where M is the mass of the kinks. The fusing tensor as linear map is bijective for the $kink + kink \to particle$ fusion.

The S-matrix $S_{PK}(\theta)$ for the scattering of a particle with $\alpha < 0$ and a kink of mass $M = -1/(2\alpha)$ is obtained from S_K by bootstrap [1, 17] applied to the $kink + kink \to particle$ vertex. It turns out that $S_{PK}(\theta)$ is also minimal and has neither poles nor zeroes in the physical strip.

It is expected that a kink is produced in the kink-particle fusion. The transformation properties of the kink-particle states discussed earlier show that in this case it is necessary that (12) is satisfied. We checked that S_{PK} is bijective (of rank eight) everywhere in the physical strip, except (12) is satisfied. In the latter case it is a projection onto the four dimensional kink subspace. The $kink + particle \to kink$ fusion is thus possible indeed, and there are no restrictions other than (12). The $kink + particle \to kink$ fusion is a crossed version of $kink + kink \to particle$ fusion. The produced kink is of the same mass as the incoming one, so the fusing angle is in the domain $[\pi/2, \pi]$. The fusing tensor is a projection.

---

[1] The same factorizations are done in the literature.



Figure 4: The two bulk vertices

Finally, there are bootstrap equations for S_P, S_K, S_{PK} and the fusing processes described above, which are expected to be valid. They were checked, and the result is [1, 3] that they are indeed satisfied: the scattering amplitudes that can be obtained from S_K, S_P and S_{PK} by bootstrap are S_K, S_P, and S_{PK} again. In particular, the condition of minimality is consistent with the bootstrap. The fusion of two particles with $\alpha < 0$ produces a particle with the same value of α. The fusion of two kinks of mass M produces a particle with $\alpha = -1/(2M)$, and if the fusing angle is ρ, then the angle u corresponding to the produced particle is $u = \pi/2 - \rho/2$. The fusion of a kink of mass M and a particle with $\alpha = -1/(2M)$ produces a kink of mass M.

The three-particle and the kink-kink-particle vertices are shown in figure 4 (the crossed versions of them are also possible). Figure 4 is the only one in which we use different line styles for different types of particles.

3 Boundary supersymmetric factorizable scattering

Let us assume that a bulk factorizable scattering theory and a supersymmetric version of this theory are given, and the supersymmetric version is of the form described in section 2.

In particular, the bulk supersymmetric theory is characterized by a single $\alpha < 0$ (if there are particles in the theory) for the scattering of any two particles, and a kink mass $M > 0$ (if there are kinks in the theory). $\alpha = -1/(2M)$ is satisfied if there are both kinks and particles in the theory. Each particle has mass $m \leq -1/\alpha = 2M$, and to each particle an angle $0 < u \leq \pi/2$ is associated as described in the previous section (see (14)). The fusing rules of the bulk theory satisfy the constraints described in section 2. These constraints are $u_{ij}^k \in \{u_i + u_j, \pi - u_i + u_j, u_i + \pi - u_j\}$ for the fusing angle of a $particle_i + particle_j \to particle_k$ process, $u_k = \pi/2 - u_{ij}^k/2$ for a $kink_i + kink_j \to particle_k$ process, and $u_{ij}^k = \pi/2 + u_i$ for a $particle_i + kink_j \to kink_k$ process.

It is also assumed that a factorizable boundary scattering theory associated to the non-supersymmetric bulk theory is also given, and the boundary is on the right hand side.

3.1 Boundary supersymmetry algebra

The presence of boundary partially violates supersymmetry. In general only one supercharge $\tilde{Q}$ and a central charge $\tilde{Z}$ is expected to be conserved if a supersymmetric boundary is present. We call the boundary version of supersymmetry boundary supersymmetry. The boundary supersymmetry algebra is denoted by $\mathcal{A}_{by}$, and the full bulk supersymmetry algebra by $\mathcal{A}_{bulk}$.

In any factorizable boundary scattering theory there is a Hilbert-space $\mathcal{H}_B$ of the boundary bound states (also containing the ground state(s)). General multi-particle *in* states containing $n \geq 1$ bulk particles traveling towards the boundary form a space $\mathcal{H}_{n,by} = \mathcal{H}_{n,b} \otimes \mathcal{H}_B$, where $\mathcal{H}_{n,b}$ is a subspace of the Hilbert-space $\mathcal{H}_n$ of bulk n-particle *in* states. $\mathcal{H}_{n,b}$ is the subspace containing *in* particles traveling towards the boundary. The Hilbert-space of the boundary theory is thus $\mathcal{H}_{by} = \mathcal{H}_B \oplus (\oplus_{n=1}^{\infty} \mathcal{H}_{n,by})$. In the supersymmetric situation a representation π_1 of $\mathcal{A}_{bulk}$ is defined on the Hilbert-space of the bulk theory, so in order to define the action of $\mathcal{A}_{by}$ on $\mathcal{H}_{by}$ we need the action π_2 of $\mathcal{A}_{by}$ on $\mathcal{H}_B$, and a coproduct $\Delta_B : \mathcal{A}_{by} \to \mathcal{A}_{bulk} \otimes \mathcal{A}_{by}$. $\mathcal{A}_{by}$ itself need not admit a coproduct. We also require the coassociativity property $(Id \otimes \Delta_B)\Delta_B = (\Delta \otimes Id)\Delta_B$, so $\mathcal{A}_{by}$ is a coideal of $\mathcal{A}_{bulk}$. A similar construction applies to the action of $\mathcal{A}_{by}$ on *out* states and *intermediate* multi-particle states. This framework is the same as that used in [15, 16] for quantum group symmetry, and it is consistent with the definition of the action of supersymmetry in [2, 8]. In fact, we need to specify the action of $\mathcal{A}_{by}$ on the ground state(s) only, because its action on the other elements of $\mathcal{H}_B$ is determined by the boundary bootstrap. We do not require in general that the action of the supersymmetry generators Q, $\bar{Q}$, and Γ should be defined on the boundary ground state(s) or on other elements of $\mathcal{H}_B$. We remark that in the construction described for instance in [2] the coproduct Δ is used, so it is necessary to specify the action of Q, $\bar{Q}$, and Γ on the ground state. $\mathcal{A}_{by}$ is a subalgebra of $\mathcal{A}_{bulk}$ in that construction, and the reflection amplitudes are invariant only with respect to $\mathcal{A}_{by}$.

In the following paragraphs we specify $\mathcal{A}_{by}$ and Δ_B. Our formulae are not derived from Lagrangian field theory in a rigorous manner, neither do they result from some rigorous classification of all coideals of $\mathcal{A}_{bulk}$, but they are consistent with considerations and results in [2, 15, 16, 7, 8], and we give further arguments supporting them. We think that the framework we describe is fairly general, and the assumptions we make are not really restrictive.

$\mathcal{A}_{by}$ is assumed to have the generators $\tilde{Q}$, $\tilde{H}$, $\tilde{Z}$ and $\tilde{I}$. $\tilde{I}$ is the identity element, $\tilde{H}$ is the time translation generator. The conservation of fermionic parity is not required, so the corresponding element $\tilde{\Gamma}$ is not included in $\mathcal{A}_{by}$. The boundary supersymmetric central charge $\tilde{Z}$ commutes with the other generators. The supercharge $\tilde{Q}$ is expected to be the boundary counterpart of Q and $\bar{Q}$, so we expect that $\tilde{Q}^2$ is of the form

$$\tilde{Q}^2 = 2\tilde{H} + p(\tilde{I}, \tilde{Z}), \tag{23}$$

where p is a polynomial. This relation implies that $\mathcal{A}_{by}$ is a commutative algebra. It is natural to assume that in analogy with the bulk SUSY algebra

$$\Delta_B(\tilde{H}) = H \otimes \tilde{I} + I \otimes \tilde{H} \quad \text{and} \quad \Delta_B(\tilde{Z}) = \hat{Z} \otimes \tilde{I} + I \otimes \tilde{Z}. \tag{24}$$

Because of the expected role of $\tilde{Q}$ and the relation (23) for $\tilde{Q}^2$ we take the Ansatz $\Delta_B(\tilde{Q}) = Q \otimes X_1 + \bar{Q} \otimes X_2 + \Gamma \otimes X_3 + I \otimes X_4 + \hat{Z} \otimes X_5$. Solving the coassociativity condition and (23) for the X_i-s we found that

$$\Delta_B(\tilde{Q}) = (Q \pm \bar{Q}) \otimes \tilde{I} + \Gamma \otimes \tilde{Q}, \tag{25}$$

$$\tilde{Q}^2 = 2\tilde{H} \pm 2\tilde{Z} + 2u\tilde{I}, \tag{26}$$

where u is an arbitrary number. Two cases can be distinguished corresponding to the sign in (25) and (26), they will be referred to as (+) and (−) cases, respectively. u can be tuned by the redefinition of $\tilde{H}$: $\tilde{H} \to \tilde{H} + u'\tilde{I}$ results in $u \to u - u'$. u can also be tuned by a similar redefinition of $\tilde{Z}$. We set u equal to zero by the redefinition $\tilde{H} + u\tilde{I} \to \tilde{H}$:

$$\tilde{Q}^2 = 2\tilde{H} \pm 2\tilde{Z}. \tag{27}$$

Another way to obtain the formula (25) is the following: let us adopt first that $\Delta_B(\tilde{Q}) = (Q \pm \bar{Q}) \otimes \tilde{I} + \textit{a boundary contribution}$, i.e. $\tilde{Q}$ is the boundary counterpart of $Q \pm \bar{Q}$. This is supported by considerations in the Lagrangian framework [2, 8, 7]. It is reasonable [2] now to adopt (27) as the boundary version of the bulk formula $(Q \pm \bar{Q})^2 = 2H \pm 2\hat{Z}$. Now $\mathcal{A}_{by}$ is isomorphic to a subalgebra $\mathcal{A}'_{by}$ of $\mathcal{A}_{bulk}$, the isomorphism i is given by $\tilde{Q} \mapsto Q \pm \bar{Q}$, $\tilde{H} \mapsto H$, $\tilde{Z} \mapsto \hat{Z}$. If $\Delta(\mathcal{A}'_{by}) \subset \mathcal{A}_{bulk} \otimes \mathcal{A}'_{by}$, which can easily be verified to be true, then we can define Δ_B using Δ as follows: $\Delta_B(X) = (Id \otimes i)^{-1}\Delta(i(X))$, $X \in \mathcal{A}_{by}$. This definition guarantees coassociativity, and it is easy to check that it leads to (25), (24). If we took (26) with nonzero u, then $i(\tilde{Q}) = Q \pm \bar{Q} + \sqrt{u}\Gamma$ would be an appropriate choice. The formulae (25), (24) are independent of u.

The boundary supersymmetry algebra as described above is isomorphic to a coideal subalgebra of $\mathcal{A}_{bulk}$. This situation and the construction above for the boundary supersymmetry algebra is very similar to that described in [15, 16] for quantum group symmetry.

$\Delta_B(\tilde{Q})$, $\Delta_B(\tilde{H})$ and also $\Delta_B(\tilde{Z})$ is a sum of two terms. The first terms can be regarded as the bulk parts of these quantities, and the second terms can be regarded as boundary contributions. This structure is in accord with the expectation that the quantities $\tilde{Q}$, $\tilde{H}$, $\tilde{Z}$ are (semi)local in the underlying Lagrangian theory. Locality also implies that the bulk and the boundary contributions commute, which is also true for our formulae (see also [2]).

If in a certain model the ground state is a singlet, then we have

$$\tilde{Q}|\,B\rangle = \gamma|\,B\rangle, \quad \tilde{Z}|\,B\rangle = z|\,B\rangle, \tag{28}$$

where $|\,B\rangle \in \mathcal{H}_B$ is the ground state, γ and z are numbers ($z \in \mathbb{R}$). For the Hamiltonian operator

$$\tilde{H}|\,B\rangle = h|\,B\rangle = ((\gamma^2/2) \mp z)|\,B\rangle$$

holds if $u = 0$. We expect that only γ is a true parameter of the model. γ is expected to be expressible in terms of the parameters of the bulk and boundary parts of the underlying classical Lagrangian density. On the subspace $(\oplus_{n=1}^{\infty}\mathcal{H}_{n,by}) \otimes |\,B\rangle$ the generators are $\tilde{Q} = Q \otimes I \pm \bar{Q} \otimes I + \gamma\Gamma \otimes I$ (cf. [2]), $\tilde{Z} = \hat{Z} \otimes I + zI \otimes I$, $\tilde{H} = H \otimes I + hI \otimes I$. (The representations π_1, π_2 are not explicitely designated.)

It is not surprising that $\mathcal{A}_{by}$ is isomorphic to a subalgebra of $\mathcal{A}_{bulk}$, because $\mathcal{A}_{by}$ acts on the singlet $|\,B\rangle$ by multiplications, so $\pi_3(\mathcal{A}_{by})$, where π_3 is the representation of $\mathcal{A}_{by}$ on $(\oplus_{n=1}^{\infty}\mathcal{H}_{n,by}) \otimes |\,B\rangle$, is a subalgebra of $\pi_2(\mathcal{A}_{bulk}) \otimes I$.

A third way to obtain (25) and (26) is the following: consider the representation of $\mathcal{A}_{by}$ on $(\oplus_{n=1}^{\infty}\mathcal{H}_{n,by}) \otimes |\,B\rangle$. Using (24) and (28) we have $\tilde{Z} = \hat{Z} \otimes I + zI \otimes I$, $\tilde{H} = H \otimes I + hI \otimes I$ on $(\oplus_{n=1}^{\infty}\mathcal{H}_{n,by}) \otimes |\,B\rangle$. Considering (23) and (28) we take the Ansatz $\tilde{Q} = aQ \otimes I + b\bar{Q} \otimes I + c\Gamma \otimes I + dI \otimes I + e\hat{Z} \otimes I$ on $(\oplus_{n=1}^{\infty}\mathcal{H}_{n,by}) \otimes |\,B\rangle$, where a, b, c, d, e are numbers. One can now solve (23) for a, b, c, d, e. The result is $\tilde{Q} = Q \otimes I \pm \bar{Q} \otimes I + \gamma\Gamma \otimes I$ and the formula (26), with $h = \gamma^2/2 \mp z - u$. The precise structure of $\mathcal{A}_{by}$ is thus obtained and the algebra is also realized as a subalgebra of $\mathcal{A}_{bulk}$. $\Delta_B(\tilde{Q})$ can be derived from Δ as earlier, or it can be found by using coassociativity and the property that Δ_B is an algebra homomorphism.

To describe situations when the fermionic parity is also conserved, $\mathcal{A}_{by}$ can be supplemented by the boundary fermionic parity generator $\tilde{\Gamma}$. It satisfies the following relations:

$$\tilde{\Gamma}^2 = \tilde{I}, \quad [\tilde{\Gamma}, \tilde{Z}] = 0, \quad [\tilde{\Gamma}, \tilde{H}] = 0, \tag{29}$$

and also

$$\{\tilde{\Gamma}, \tilde{Q}\} = 2g\tilde{I}, \tag{30}$$

where g is a parameter. The coproduct of $\tilde{\Gamma}$ is

$$\Delta_B(\tilde{\Gamma}) = \Gamma \otimes \tilde{\Gamma}. \tag{31}$$

Formula (30) can be obtained from the requirement that Δ_B is a homomorphism.

In case of a model with singlet boundary ground state

$$\tilde{\Gamma}|\,B\rangle = \epsilon|\,B\rangle, \qquad \epsilon = \pm 1,$$

and $g = \epsilon\gamma$. The sign ϵ is not expected to be a true parameter of the model (see also [2]).

3.2 Supersymmetrized reflection amplitudes

We assume that —similarly to the bulk case— the boundary states $|\,\eta_i\rangle$ of the non-supersymmetric theory become multiplets $|\,B_i\rangle \otimes |\,\eta_i\rangle$ in the supersymmetrized boundary theory, and that the one-particle reflection amplitude $\hat{R}$ on these states take the form

$$\hat{R}^{\xi'_i A'_i;\eta'_j B'_j}_{\xi_i A_i;\eta_j B_j}(\theta) = R^{A'_i B'_j}_{A_i B_j}(\theta)\tilde{R}^{\xi'_i \eta'_j}_{\xi_i \eta_j}(\theta), \tag{32}$$

where $\tilde{R}$ is the one-particle reflection amplitude in the non-supersymmetric model, and R is called the supersymmetric factor. B_i specify the boundary supersymmetric quantum numbers, boundary supersymmetry acts trivially on the non-supersymmetric indices. The index i specifies quantum numbers which are conserved in the non-supersymmetric boundary theory. These quantum numbers should be defined and conserved in the bulk theory as well, and in the bulk theory they should be Poincare and charge conjugation invariant. This implies that the bulk SUSY representations in which the particles transform should depend only on quantum numbers which are also conserved in reflections. This is a restriction on the bulk supersymmetric theory in principle. It is also required that the reflection amplitude $\hat{R}$ and so also R commute with the boundary supersymmetry.

The factorized form of $\hat{R}$ (and $\hat{S}$) implies that the boundary Yang-Baxter equations, the boundary unitarity equation and the boundary cross-unitarity equation also factorize, so R has to satisfy them separately in order for $\hat{R}$ to satisfy them. The boundary fusing equations and fusing tensors are also assumed to factorize, and this implies that the boundary bootstrap equations also factorize. The non-supersymmetric part of the bootstrap equations are satisfied, so we require that the bootstrap equations should be satisfied by the supersymmetric factors separately.

As in the bulk case, the non-supersymmetric boundary (and bulk) fusing data enter the fusing and bootstrap equations for the supersymmetric factors, and this places nontrivial conditions on the described supersymmetrization scheme.

It is required that the supersymmetric factors of the reflection amplitudes should be minimal, without poles and overall zeroes on the imaginary axis in the "physical strip", which is $0 < Im(\theta) < \pi/2$ for the rapidity argument θ of the reflection amplitudes. This guarantees that no new bound states are introduced by the supersymmetrization.

The supersymmetric boundary fusing equations are analogous with the bulk ones:

$$R^{A'_i B'_j}_{A_i B_j}(i\nu^k_{ij}) = \sum_{B_k} g^{B_k}_{A_i B_j} h^{A'_i B'_j}_{B_k}, \qquad |\, A_i B_j\rangle = \sum_{B_k} g^{B_k}_{A_i B_j} |\, B_k\rangle, \quad |\, B_k\rangle = \sum_{A_i, B_j} h^{A_i B_j}_{B_k} |\, A_i B_j\rangle,$$

ν^k_{ij} is the boundary fusing angle for the $\xi_i + \eta_j \to \eta_k$ fusion, g and h are the boundary fusing and decay tensors respectively. For graphical representation see fig. 5.

3.3 Ground state representation

There are several different representations of the boundary supersymmetry algebra which can serve as representations in which the ground states of various theories transform. Moreover, it is not necessarily a natural assumption that the supersymmetric part of the



Figure 5: Boundary fusing equation (splitting of R)

ground state of a certain theory is a singlet. Nevertheless, in this paper we consider theories which can be supersymmetrized in such a way that the ground state is a singlet with RSOS label $\frac{1}{2}$ only. In particular, the corresponding adjacency condition between the ground state and the nearest kink is required to be satisfied. The supersymmetric part of the ground state is denoted by $|\,B_{\frac{1}{2}}\rangle$. The action of the boundary supersymmetry generators in this case on $|\,B_{\frac{1}{2}}\rangle$ is

$$\tilde{Q}^{(-)}|\,B_{\frac{1}{2}}\rangle = e\gamma|\,B_{\frac{1}{2}}\rangle, \quad \tilde{Z}|\,B_{\frac{1}{2}}\rangle = 0, \quad e = \pm 1, \tag{33}$$

where

$$\gamma \in \mathbb{R}, \quad \gamma < 0, \qquad \text{or} \qquad \gamma \in i\mathbb{R}, \quad \gamma/i < 0,$$

and

$$\tilde{Q}^{(+)}|\,B_{\frac{1}{2}}\rangle = \gamma|\,B_{\frac{1}{2}}\rangle, \quad \tilde{Z}|\,B_{\frac{1}{2}}\rangle = 0,$$

where

$$\gamma \in \mathbb{R}, \qquad \text{or} \qquad \gamma \in i\mathbb{R}.$$

The superscript of $\tilde{Q}$ stands for the $(+)$ and $(-)$ cases. $e\gamma$ or γ is a parameter of the model to be described and is expected to be expressible in terms of the parameters of the Lagrangian density. The reason for writing this parameter in the form $e\gamma$ in the $(-)$ case will become clear in 3.5 when the ground state kink reflection amplitudes are discussed. Two (symmetric) cases correspond to the sign e. We remark that singlet ground states with RSOS label 1 or 0 are also possible.

3.4 Factorized scattering

Let the supersymmetric factor of the amplitude of a general multi-particle process be A. The rapidities of the particles are allowed to take complex values. Assume that there are m individual two-particle scatterings, reflections, fusions or decays in the process, which are labeled by $1, 2, ..., m$ in chronological order ($t_1 < t_2 < ... < t_m$). A can be written as a product of m tensors, which can also be viewed as linear maps:

$$A = T_m T_{m-1} T_{m-2} ... T_k ... T_1,$$

where the multiplication is the multiplication of linear maps, i.e. the contraction of the appropriate upper and lower indices. If at the time t_k the "event" is the scattering of the n_k-th and (n_k+1)-th particle and by this time the *intermediate* state contains m_k one-particle states, then

$$T_k = I_1 \otimes I_2 \otimes ... \otimes I_{n_k-1} \otimes S_{n_k,n_k+1} \otimes I_{n_k+2} \otimes ... \otimes I_{m_k},$$

where S_{n_k,n_k+1} is the two-particle S-matrix for the n_k-th and (n_k+1)-th particle. The I_l is the identity tensor for the SUSY indices of the l-th particle or for the SUSY indices of the boundary state. Similar formulae can be written if the "event" at t_k is a reflection, fusion or decay.

The factorized form above has (among others) the following simple implications: if S_{n_k,n_k+1} (or the reflection amplitude, fusing or decay tensor) is bijective as a linear map, then T_k is also bijective. If all the $T_1,...T_m$-s are bijective, then so is A. The T_k-s are equivariant.

3.5 Supersymmetric reflection factors on the ground state boundary

We consider one-particle kink reflection factors on ground state boundary first. As the left and right RSOS labels should be conserved, $\{R_K\}_{K_{1\frac{1}{2}}}^{K_{0\frac{1}{2}}}(\theta) = \{R_K\}_{K_{0\frac{1}{2}}}^{K_{1\frac{1}{2}}}(\theta) = 0$ must hold, i.e. R_K is diagonal (R_K denotes the kink reflection factor). The general solution of the boundary Yang-Baxter equation, unitarity condition and crossing equation without imposing supersymmetry is [14, 9]

$$\{R_K\}_{K_{0\frac{1}{2}}}^{K_{0\frac{1}{2}}}(\theta) = (1 + A\sinh(\theta/2))M(\theta), \qquad \{R_K\}_{K_{1\frac{1}{2}}}^{K_{1\frac{1}{2}}}(\theta) = (1 - A\sinh(\theta/2))M(\theta),$$

where $M(\theta)$ is restricted by unitarity and crossing symmetry. After imposing the boundary supersymmetry condition one finds that in the $(+)$ case [2, 8]

$$\{R_K^{(+)}\}_{K_{0\frac{1}{2}}}^{K_{0\frac{1}{2}}}(\theta) = \{R_K^{(+)}\}_{K_{1\frac{1}{2}}}^{K_{1\frac{1}{2}}}(\theta) = 2^{-\theta/(\pi i)}P(\theta), \qquad A = 0.$$

($P(\theta)$ can be found in the Appendix.) In the $(-)$ case there are distinct solutions for a given γ corresponding to the sign e:

$$\{R_{K,e}^{(-)}\}_{K_{0\frac{1}{2}}}^{K_{0\frac{1}{2}}}(\theta) = (\cos\frac{\xi}{2} + ei\sinh\frac{\theta}{2})K(\theta - i\xi)K(i\pi - \theta - i\xi)2^{-\theta/(\pi i)}P(\theta), \qquad (34)$$

$$\{R_{K,e}^{(-)}\}_{K_{1\frac{1}{2}}}^{K_{1\frac{1}{2}}}(\theta) = (\cos\frac{\xi}{2} - ei\sinh\frac{\theta}{2})K(\theta - i\xi)K(i\pi - \theta - i\xi)2^{-\theta/(\pi i)}P(\theta), \qquad (35)$$

where $\gamma = -2\sqrt{M}\cos\frac{\xi}{2}$ and $0 \leq \xi \leq \pi$ or $Re(\xi) = 0$ or $Re(\xi) = \pi$, M is the kink mass. (34) and (35) are invariant under $\xi \leftrightarrow -\xi$. These reflection amplitudes are minimal, they have no poles and zeroes in the physical strip. The sign e seems to correspond here to the 0 and 1 RSOS vacua. It should be noted that $R_K^{(+)}(\theta)$ is independent of γ and e. Furthermore, as symmetry under $\tilde{\Gamma}$ requires $R_{K_{0\frac{1}{2}}}^{K_{0\frac{1}{2}}}(\theta) = R_{K_{1\frac{1}{2}}}^{K_{1\frac{1}{2}}}(\theta)$, the $R_K^{(+)}(\theta)$ is automatically $\tilde{\Gamma}$-symmetric, although this is not required a priori. On the other hand, $R_{K,e}^{(-)}(\theta)$ are not $\tilde{\Gamma}$-symmetric. However, $\{R_{K,e}^{(-)}\}_{K_{0\frac{1}{2}}}^{K_{0\frac{1}{2}}}(\theta) = -\{R_{K,e}^{(-)}\}_{K_{1\frac{1}{2}}}^{K_{1\frac{1}{2}}}(\theta)$ if $\gamma = 0$ ($A \to \infty$). We also remark that $\{R_K\}_{K_{1\frac{1}{2}}}^{K_{1\frac{1}{2}}}(\theta)/\{R_K\}_{K_{0\frac{1}{2}}}^{K_{0\frac{1}{2}}}(\theta)$ is determined by the supersymmetry condition, i.e. if we impose the condition of invariance under supersymmetry, then we do not need to solve the Yang-Baxter equation.

We determined the general solution of the Yang-Baxter equation for the boundary supersymmetric particle reflection (on the ground state boundary). We imposed the supersymmetry condition first, which is $Q(-\theta)R(\theta) = R(\theta)Q(\theta)$, where $Q(\theta)$ is $\tilde{Q}$ on the states of the form $\mid A(\theta)B_{\frac{1}{2}}\rangle$, and $R(\theta)$ is the particle reflection amplitude on ground state boundary. The resulting forms of the reflection amplitude in the $(+)$ and $(-)$ cases are

$$R_P^{(+)}(\theta) = Z^{(+)}(\theta)\frac{1}{\sqrt{m}}\begin{pmatrix} (X^{(+)}(\theta) + e\gamma Y^{(+)}(\theta))c(\frac{\theta}{2} - \frac{i\pi}{4}) & \sqrt{m}Y^{(+)}(\theta)c(\theta) \\ \sqrt{m}Y^{(+)}(\theta)c(\theta) & (X^{(+)}(\theta) - e\gamma Y^{(+)}(\theta))c(\frac{\theta}{2} + \frac{i\pi}{4}) \end{pmatrix},$$

$$R_P^{(-)}(\theta) = Z^{(-)}(\theta)\frac{1}{\sqrt{m}}\begin{pmatrix} (X^{(-)}(\theta) + e\gamma Y^{(-)}(\theta))c(\frac{\theta}{2} + \frac{i\pi}{4}) & i\sqrt{m}Y^{(-)}(\theta)c(\theta) \\ -i\sqrt{m}Y^{(-)}(\theta)c(\theta) & (X^{(-)}(\theta) - e\gamma Y^{(-)}(\theta))c(\frac{\theta}{2} - \frac{i\pi}{4}) \end{pmatrix},$$

where c stands for cosh and X, Y and Z are functions not determined by supersymmetry. Now two cases can be distinguished depending on whether $\tilde{\Gamma}$ is a symmetry or not: in the first case, which is the $\tilde{\Gamma}$-symmetric case, $Y(\theta) \equiv 0$, $X(\theta)$ can be absorbed into the prefactor, and the structure of the reflection amplitude is completely determined and does not contain free parameters:

$$R_{P1}^{(\pm)}(\theta) = \frac{1}{\sqrt{m}}ZX^{(\pm)}(\theta)\begin{pmatrix} \cosh(\frac{\theta}{2} \mp \frac{i\pi}{4}) & 0 \\ 0 & \cosh(\frac{\theta}{2} \pm \frac{i\pi}{4}) \end{pmatrix}.$$

This case is discussed in [4], the explicit form of $ZX^{(\pm)}$ can be found in the Appendix, see also [4, 10, 2]. It can be checked that the boundary Yang-Baxter equation for incoming particles of arbitrary masses is satisfied by this reflection amplitude. $R_{P1}^{(\pm)}(\theta)$ can also be obtained from $R_K^{(+)}(\theta)$ and $R_K^{(-)}(\theta)$ at $\gamma = 0$ by bootstrap [2, 10].

In the second case, when $\tilde{\Gamma}$ is not conserved, $Y(\theta)$ is not identically zero, and it can be absorbed into the prefactor, so one free function $y^{(\pm)}(\theta) = X^{(\pm)}(\theta)/Y^{(\pm)}(\theta)$ remains in the reflection amplitude, which is to be determined by the boundary Yang-Baxter equation. To obtain $y^{(\pm)}(\theta)$ we solved the Yang-Baxter equation first in the case when the conserved quantum numbers have the same values for the two incoming particles.

Although the boundary Yang-Baxter equation is quadratic in general, in this case it is inhomogeneous linear in the variables $y^{(\pm)}(\theta_1)$ and $y^{(\pm)}(\theta_2)$. The coefficient of the quadratic term $y^{(\pm)}(\theta_1)y^{(\pm)}(\theta_2)$ vanishes precisely because $R^{(\pm)}_{P1}(\theta)$ (!) satisfies the Yang-Baxter equation. The Yang Baxter equation consists of 16 scalar equations in our case. Some of them are trivial (0=0), and the remaining n equations are of the form

$$a^{(\pm)}_q(\theta_1,\theta_2)y^{(\pm)}(\theta_1)+b^{(\pm)}_q(\theta_1,\theta_2)y^{(\pm)}(\theta_2)+c^{(\pm)}_q(\theta_1,\theta_2)=0,\quad q=1..n.$$

It is possible to choose two inequivalent equations from this set. Two such equations can be solved for the numbers $y^{(\pm)}(\theta_1)$ and $y^{(\pm)}(\theta_2)$. The solution turns out to be of the form $y^{(\pm)}(\theta_1)=g^{(\pm)}(\theta_1)$, $y^{(\pm)}(\theta_2)=g^{(\pm)}(\theta_2)$ (for general coefficients $d_{q_1}, e_{q_1}, f_{q_1}, d_{q2}, e_{q2}, f_{q2}$ it would be of the form $y(\theta_1)=g_1(\theta_1,\theta_2)$, $y(\theta_2)=g_2(\theta_1,\theta_2)$, which does not define a function $y(\theta)$), where $g^{(\pm)}$ is a function that depends also on m, α, γ, but has no other parameters. Consequently, the reflection amplitude depends on the conserved quantum numbers through these parameters only. We checked that the solution obtained in this way satisfies the other $n-2$ equations as well. In the next step we checked that the solutions $R^{(\pm)}_{P2,e}(\theta)$ satisfy the Yang-Baxter equation for incoming particles of different masses as well. The two functions $y^{(+)}(\theta)$ and $y^{(-)}(\theta)$ have very similar form.

The solutions that we obtained can be brought to the following form:

$$\{R^{(\pm)}_{P2,e}\}^b_b(\theta)=A^{(\pm)}_+(\theta),\quad \{R^{(\pm)}_{P2,e}\}^f_f(\theta)=A^{(\pm)}_-(\theta),$$

$$\{R^{(\pm)}_{P2,e}\}^f_b(\theta)=\pm B^{(\pm)}(\theta),\quad \{R^{(\pm)}_{P2,e}\}^b_f(\theta)=B^{(\pm)}(\theta),$$

$$\begin{aligned}A^{(-)}_\pm(\theta)=\tilde{Z}^{(-)}(\theta)&\left\{\cosh(\frac{\theta}{2})\left(\frac{\gamma^2}{4M}-\left[\sin^2(\frac{\rho}{4})+\sinh^2(\frac{\theta}{2})\right]\right)\right.\\&\left.\mp i\sinh(\frac{\theta}{2})\left(\frac{\gamma^2}{4M}+\left[\sin^2(\frac{\rho}{4})+\sinh^2(\frac{\theta}{2})\right]\right)\right\},\\A^{(+)}_\pm(\theta)=\tilde{Z}^{(+)}(\theta)&\left\{-i\cosh(\frac{\theta}{2})\left(\frac{\gamma^2}{4M}-\left[\sin^2(\frac{\rho}{4})-\cosh^2(\frac{\theta}{2})\right]\right)\right.\\&\left.\pm\sinh(\frac{\theta}{2})\left(\frac{\gamma^2}{4M}+\left[\sin^2(\frac{\rho}{4})-\cosh^2(\frac{\theta}{2})\right]\right)\right\}\\B^{(\pm)}(\theta)=\tilde{Z}^{(\pm)}(\theta)&\frac{e\gamma}{2\sqrt{M}}\sqrt{\cos(\rho/2)}\sinh(\theta),\end{aligned}$$

where

$$m=2M\cos(\frac{\rho}{2}),\quad \frac{\rho}{2}=\frac{\pi}{2}-u,\quad \alpha=-\frac{1}{2M},\tag{36}$$

$0\le\rho<\pi$, $e=\pm1$ in the $(-)$ case and $e=1$ in the $(+)$ case. Note that $R^{(\pm)}_{P2,e}$ depends on two parameters: γ^2/M and ρ only. $R^{(-)}_{P2,e}$ has the same structure as the particle reflection amplitude obtained in [2] for the case of the boundary supersymmetric sine-Gordon model from the kink reflection amplitude by bootstrap. Consequently, there is no need now to solve the crossing and unitarity equations for $\tilde{Z}^{(-)}(\theta)$, we take it from [2]. We determined $\tilde{Z}^{(+)}(\theta)$ using the unitarity and crossing equations and exploiting the fact that these equations take a similar form for $\tilde{Z}^{(-)}(\theta)$. Explicit formulae for these prefactors can be found in the Appendix.

In the $(+)$ case we introduce the parameter ξ in the following way: $\gamma = -2\sqrt{M} i \sin(\xi/2)$, $-\pi \leq \xi \leq \pi$ or $Re(\xi) = 0$ or $Re(\xi) = \pm\pi$.

To summarize, we have the solutions $R^{(\pm)}_{P2,e}$, which depend on γ and e and are not symmetric with respect to $\tilde{\Gamma}$, and we also have $R^{(\pm)}_{P1}$, which are independent of γ, e and are symmetric with respect to $\tilde{\Gamma}$. $R^{(\pm)}_{P1}$ and $R^{(\pm)}_{P2,e}$ do not satisfy the Yang-Baxter equation together. It is also important to note that $\lim_{\gamma\to 0} R^{(\pm)}_{P2,e} = R^{(\pm)}_{P1}$. The poles and degeneracies of these reflection amplitudes will be discussed in section 4.

The same set of kink and particle reflection factors can be obtained by solving the Yang-Baxter equations without imposing the boundary supersymmetry condition [14, 10, 9]. The supersymmetry condition relates the parameters of the reflection factors obtained in this way to the parameters of the representations of the supersymmetry algebra. The results described above show that the task of solving the Yang-Baxter equations is greatly simplified if one imposes the supersymmetry condition first.

4 Supersymmetric boundary bootstrap

The first boundary bootstrap equation (see fig. 6) applied to the $kink + kink \to particle$ bulk fusion determines reflection factors for particles on ground state boundary. They turn out [10, 2] to be the same as those obtainable by solving the boundary Yang-Baxter, crossing and unitarity equations. In terms of the amplitudes $R^{(+)}_K + R^{(+)}_K \to R^{(+)}_{P1}$ and $R^{(-)}_{K,e} + R^{(-)}_{K,e} \to R^{(-)}_{P2,e}$ (with appropriate values of the parameters). Similarly, it can be checked that the first boundary bootstrap equation is also satisfied for the $particle + particle \to particle$, $kink + particle \to kink$ fusions with the amplitudes $R^{(\pm)}_{P1} + R^{(\pm)}_{P1} \to R^{(\pm)}_{P1}$, $R^{(\pm)}_{P2,e} + R^{(\pm)}_{P2,e} \to R^{(\pm)}_{P2,e}$ and $R^{(+)}_K + R^{(+)}_{P1} \to R^{(+)}_K$, $R^{(-)}_{K,e} + R^{(-)}_{P2,e} \to R^{(-)}_{K,e}$ respectively. These relations are nontrivial, although it is clear that they are satisfied up to CDD factors. It is remarkable that $R^{(+)}_{P2,e}$ cannot be obtained by bootstrap from kink reflection factors, while the other particle reflection factors $R^{(\pm)}_{P1}$ and $R^{(-)}_{P2,e}$ can.

4.1 Properties of the ground state reflection factors

$R^{(+)}_K$ is bijective (of rank two) in the physical strip. $\{R^{(-)}_{K,+1}\}^{K_{0\frac{1}{2}}}_{K_{0\frac{1}{2}}}(\theta)$ and $\{R^{(-)}_{K,-1}\}^{K_{1\frac{1}{2}}}_{K_{1\frac{1}{2}}}(\theta)$ have a zero at $\theta = i(\pi - \xi)$, so $R^{(-)}_{K,e}$ is of rank one at this angle. This zero is in the physical strip if $\pi > \xi > \pi/2$, any other zeroes of the kink amplitudes are outside the physical strip.

Consequently, the relations $R_K^{(+)} + R_K^{(+)} \to R_{P1}^{(+)}$, and $R_{K,e}^{(-)} + R_{K,e}^{(-)} \to R_{P2,e}^{(-)}$ together with the bijectivity of the kink-kink fusing tensor and S_K imply that $R_{P1}^{(+)}$ is of rank two (bijective) and has no poles in the physical strip, and $R_{P2,e}^{(-)}(\theta)$ is also bijective for generic values of θ, but it is of rank one if $\theta = i(\pi - \xi \pm \rho/2)$. It is possible for these angles to be in the physical strip and on the imaginary axis if $Im(\xi) = 0$. In this case $(\pi - \xi - \rho/2) > -\pi/2$, so if $(\pi - \xi - \rho/2)$ is negative, then there is a pole in the physical strip at $i(\rho/2 + \xi - \pi)$ because of unitarity. If $(\pi - \xi - \rho/2) > 0$, then $R_{P2,e}^{(-)}(\theta)$ has no poles and zeroes in the physical strip, and within the physical strip it is of rank 1 if and only if $\theta = i(\pi - \xi \pm \rho/2)$. We therefore impose the following condition on ξ (and also on γ) provided that $Im(\xi) = 0$:

$$\pi - \xi \geq \rho/2. \tag{37}$$

$R_{P1}^{(-)}$ is also bijective and has no poles in the physical strip. It can be verified by direct calculation that $R_{P2,e}^{(+)}(\theta)$ is degenerate (of rank two) at $\theta = i(\pi - \xi \pm \rho/2)$ and $\theta = i(\pi + \xi \pm \rho/2)$, so the condition (37) reads in this case as

$$\pi - |\xi| \geq \rho/2. \tag{38}$$

(Note that the relation between ξ and γ is different in the $(+)$ and $(-)$ cases). If ξ is not real, then all the kink and particle reflection factors are nondegenerate on the imaginary axis within the physical strip.

The relations above imply that it is consistent with the first boundary bootstrap equation to describe the reflections on ground state boundary of all kinks and particles in a theory by the minimal $R_K^{(+)}$, $R_{K,e}^{(-)}$, $R_{P1}^{(\pm)}$, $R_{P2,e}^{(\pm)}$ solutions, which have the property that they have neither poles nor (overall) zeroes on the imaginary axis in the physical strip, so the pole structure of the (full) ground state reflection factors is the same as in the non-supersymmetric theory.

A particular boundary scattering theory is characterized apart from the parameters of the bulk theory by the sign $(+)$ or $(-)$, and also by the parameters γ and e. In the $(+)$ case the amplitudes $R_K^{(+)}$ and $R_{P1}^{(+)}$ are to be used if there are both kinks and particles in the theory. If there are only particles in the theory, then either $R_{P2,e}^{(+)}$ should be used for all particles, or $R_{P1}^{(+)}$. In the $(-)$ case the amplitudes $R_{K,e}^{(-)}$ and $R_{P2,e}^{(-)}$ should be used as ground state reflection factors if there are both kinks and particles in the theory. If there are only particles in the theory, then either $R_{P2,e}^{(-)}$ or $R_{P1}^{(-)}$ should be used for all particles.

(37) and (38) imply that if $R_{P2,e}^{(\pm)}$ describes the ground state reflections of the particles in a theory, then

$$\pi - |\xi| \geq \rho_{max}/2, \qquad \rho_{max} = \max_i(\rho_i), \tag{39}$$



Figure 6: Boundary bootstrap equation I (BBE I)

where i runs over all particles, is necessary and sufficient for all $R^{(\pm)}_{P2,e}$ to have no poles in the physical strip. We remark, that the following CDD factors are consistent with the BBE I: $E(\theta) = C(\theta - i\rho/2)C(\theta + i\rho/2)$ for particles with mass $2M\cos(\rho/2)$, and $C(\theta)$ for kinks, where $C(\theta)$ is an arbitrary CDD factor.

We introduce the following notation for the supersymmetric part of boundary bound states: $|\,\nu_1, \nu_2, ..., \nu_k, B_{\frac{1}{2}}\rangle$ denotes a multiplet (subspace) of states, where ν_1, ν_2,...,ν_k are the fusing angles in the successive steps of the creation of the multiplet of states. The incoming bulk particle multiplet in the nth step of the creation is $p_n(i\nu_n)$. It is either a kink or a boson/fermion multiplet, and if it is a boson/fermion multiplet, then also an angle u_n is associated to it (see (14)). We will sometimes use the letter μ instead of ν if the incoming bulk particle multiplet is a kink multiplet. We usually use the term "state" for a multiplet (subspace) of states from now on. We call two states $|\,\nu_1, \nu_2, ..., \nu_k, B_{\frac{1}{2}}\rangle$ and $|\,\nu'_1, \nu'_2, ..., \nu'_{k'}, B_{\frac{1}{2}}\rangle$ equal if and only if $k = k'$, $\nu_i = \nu'_i$, and the type (boson/fermion or kink) and mass of p_i is the same as that of p'_i $\forall i = 1...k$. In other words, two states are regarded equal if they are generated in the same way. The number $k_1 + k_2/2$ will be called the level of the boundary state, where k_1 is the number of particles and k_2 is the number of kinks among the $k = k_1 + k_2$ incoming particles. The notation for several kinks or particles in the bulk is $|\,p(i\theta_1), ..., p(i\theta_n), \nu_1, ..., \nu_k, B_{\frac{1}{2}}\rangle$ if the boundary is in the state $|\,\nu_1, ..., \nu_k, B_{\frac{1}{2}}\rangle$. $i\theta_1 ... i\theta_n$ are the rapidities and $k(i\theta)$ is also used instead of $p(i\theta)$ for kinks. The "$B_{\frac{1}{2}}$" will often be omitted.

Two states will be called equivalent if they can be mapped to each other by a bijective supersymmetry-equivariant map so that their corresponding reflection and scattering amplitudes on any other states are also equal. Our aim is to characterize the supersymmetric parts of the states up to equivalence.

We introduce the shorthand $p_1(i\theta_1)Fp_2(i\theta_2)$ for the statement "$p_1(i\theta_1)$ and $p_2(i\theta_2)$ satisfy the bulk fusing conditions described in section 2". For the converse of this statement we use the shorthand $p_1(i\theta_1)nFp_2(i\theta_2)$.



Figure 7: Boundary bootstrap equation II (BBE II)

4.2 Supersymmetric boundary states generated by particles

In this and subsequent sections the subscripts and superscripts introduced for two-particle S-matrices and one-particle reflection matrices will often be suppressed, but this should not cause any confusion. The following discussion applies to all the cases described in the previous section. In this section and in 4.3 we assume, that the ground state kink or particle reflection factors used are nondegenerate at the rapidities $i\nu_1, ..., i\nu_k$, $|\,\nu_1, ..., \nu_k\rangle$ being the state under consideration. The discussion of the case when this condition is not satisfied is deferred until the end of section 4.4.

The following arguments include diagrams of processes, states, etc., but it is important that they are to be understood to represent algebraic objects, expressions and relations. It is not required that the particular non-supersymmetric theory to be supersymmetrized contains the counterparts of all the states (for example) that will be introduced.

4.2.1 First level boundary states

A first level boundary bound state $|\,\nu\rangle$ has multiplicity 2, and the corresponding fusing tensors are bijective. Reflection amplitudes on a first level boundary state $|\,\nu\rangle$ are obtained by the second bootstrap equation (see fig. 7):

$$R_1(\theta)[I \otimes g] = [I \otimes g][S_P(\theta + i\nu) \otimes I][I \otimes R(\theta)][S_P(\theta - i\nu) \otimes I], \qquad (40)$$

where θ is the rapidity of the reflecting particle p_1, R is a ground state particle reflection amplitude, g is a boundary fusing tensor, and R_1 is the reflection amplitude to be determined.

It is clear from the r.h.s. of eq. (40) that $R_1(\theta)$ has no poles if $\theta - i\nu$ (and also θ) is on the imaginary axis in the physical strip. This does not guarantee that $R_1(\theta)$ has no poles in the physical strip (on the imaginary axis) at all, because $\theta - i\nu$ is not necessarily in the physical strip. $S_P(\theta - i\nu)$ on the r.h.s. has a pole at θ precisely if $p(i\nu)Fp_1(\theta)$, where $\nu > \theta/i$. In this case R_1 also has a pole at θ. It can easily be checked that if $p(i\nu)Fp_1(\theta)$ then $S_P(\theta + i\nu)$ is bijective.

Consequently, assuming that $|\,\nu\rangle$ has multiplicity 2, $R_1(\theta)$ has no poles on the imagi-



Figure 8:

nary axis in the physical strip if and only if $p(i\nu)Fp_1(\theta)$ is impossible for $\theta/i < \nu$, i.e.

$$\nu \leq u + u_1, \tag{41}$$

where u is the angle associated (see (14)) to the particle that generates the boundary state $|\,\nu\rangle$ and u_1 is the angle associated to the reflecting particle $p_1(\theta)$. This implies that in order for the supersymmetric reflection amplitudes of all particles in a theory on the boundary state $|\,\nu\rangle$ to have no poles on the imaginary axis in the physical strip it is necessary and sufficient that

$$\nu \leq u + u_{min}, \tag{42}$$

where $u_{min} = \min_i u_i$, i runs over all particles in the theory.

4.2.2 The multiplicity of second level boundary states

Let $|\,\nu_1, \nu_2\rangle$ be a boundary bound state and assume that $|\,\nu_1\rangle$ satisfies (42). We consider first the case when $\nu_2 > \nu_1$. In this case the amplitude A shown on the l.h.s of figure 8 can be used to determine the multiplicity and transformation properties of $|\,\nu_1, \nu_2\rangle$. The following equations hold in this case (see also fig. 8):

$$[I \otimes h_1]h_2 g_2[I \otimes g_1] = [I \otimes h_1]R_{21}(i\nu_2)[I \otimes g_1] =$$

$$\begin{aligned} &= [I \otimes h_1][I \otimes g_1][S_P(i\nu_2 + i\nu_1) \otimes I][I \otimes R(i\nu_2)][S_P(i\nu_2 - i\nu_1)I] = \\ &= [I \otimes R(i\nu_1)][S_P(i\nu_1 + i\nu_2) \otimes I][I \otimes R(i\nu_2)][S_P(i\nu_2 - i\nu_1) \otimes I]. \end{aligned} \tag{43}$$

If $S_P(i\nu_1 + i\nu_2)$ and $S_P(i\nu_1 - i\nu_2)$ are bijective, then the multiplicity of the state $|\,\nu_1, \nu_2\rangle$ is 4, and it transforms as the two-particle state $|\,p(i\nu_2), p(i\nu_1), B_{\frac{1}{2}}\rangle$.

If $S_P(i\nu_1 - i\nu_2)$ is degenerate, then the equation represented in fig. 9a. holds.

It is not hard to see, that the ground state reflection factor R shown on the figure is bijective, so the equation implies that the multiplicity of the state $|\,\nu_1, \nu_2\rangle$ is 2 and it is equivalent to $|\,p(i\nu'), B_{\frac{1}{2}}\rangle$. Continuing the equation shown on figure 9a. one can introduce a new first level state $|\,\nu'\rangle$. This is shown on figure 9b. $|\,\nu'\rangle$ is equivalent to $|\,\nu_1, \nu_2\rangle$. It is easy to verify that ν' also satisfies condition (42).

If $S_P(i\nu_1 - i\nu_2)$ is bijective but $S_P(i\nu_1 + i\nu_2)$ is not, then the equation corresponding either to fig. 11a. or to fig. 11b. holds, and it shows that the multiplicity of the state



(a) (b)

Figure 9:



Figure 10:

$|\,\nu_1,\nu_2\rangle$ is 2 and it is equivalent to $|\,p(i\nu'),B_{\frac{1}{2}}\rangle$. ($R$ is bijective.) Again a new $|\,\nu''\rangle$ first level state can be introduced, which is equivalent to $|\,\nu_1,\nu_2\rangle$. It is easy to verify that ν'' also satisfies condition (42).

In case of $\nu_2<\nu_1$ we consider the amplitude shown on the left hand side of figure 10. The condition (42) implies that $S_P(i\nu_1-i\nu_2)$ is bijective, so this amplitude is useful for determining the multiplicity of $|\,\nu_1,\nu_2\rangle$. An equation (which is analogous to (43)) corresponding to figure 10 is satisfied, and the same considerations apply to the r.h.s. as in the $\nu_1<\nu_2$ case. It is clear in particular that $|\,\nu_1,\nu_2\rangle$ is equivalent to $|\,\nu_2,\nu_1\rangle$ if these states have multiplicity 4, i.e. if $S_P(i|\nu_1-\nu_2|)$ and $S_P(i\nu_1+i\nu_2)$ are bijective.



(a) (b)

Figure 11:

4.2.3 Scattering on second level boundary states

Let $|\,\nu_1,\nu_2\rangle$ be a second level boundary bound state for which $S_P(i|\nu_1-\nu_2|)$ and $S_P(i\nu_1+i\nu_2)$ are bijective. In any other case $|\,\nu_1,\nu_2\rangle$ is equivalent to some lower level state, and scattering on $|\,\nu_1,\nu_2\rangle$ is equivalent to scattering on that lower level state. The same argument as in subsection 4.2.1 gives that condition (42) has to be satisfied also by ν_2 in order for the reflection amplitude on $|\,\nu_1,\nu_2\rangle$ to have no poles on the imaginary axis in the physical strip.

4.2.4 Boundary states of arbitrary level

The following results can be obtained by applying for higher level states essentially the same steps and arguments as described in the previous subsections. The strategy is the following: a multi-particle reflection amplitude like the one on the l.h.s. of figure 8 or 10 should be considered at the values of rapidities corresponding to the boundary bound state under consideration. The corresponding diagram should be transformed into a diagram in which no lines corresponding to boundary states occur (as on fig. 8). Then the vertices at which the corresponding two-particle scattering amplitudes are degenerate should be split, and the fusing and decaying points should be moved sufficiently far away in the future and in the past, while keeping the rest of the diagram fixed. If more than one vertices can be split, then one of them should be chosen appropriately. Finally, if there are no more vertices to split, then the diagram is in a form that shows the transformation and scattering properties of the boundary state manifestly.

Proposition 1: A general boundary bound state is equivalent to a boundary state $|\,\nu_1,\nu_2,...,\nu_n\rangle$, which is such that ν_i satisfies condition (42) for $i=1..n$, and $p(i\nu_i)nFp(i\nu_j)$ and
$p(i\nu_i)nFp(-i\nu_j)$ are satisfied for any $i,j=1..n$, $i\neq j$. The multiplicity of this state is 2^n and the state is equivalent to $|\,p(i\nu_1),p(i\nu_2),...,p(i\nu_n),B_{\frac{1}{2}}\rangle$. The reflection amplitudes on the state $|\,\nu_1,\nu_2,...,\nu_n\rangle$ have no poles on the imaginary axis in the physical strip. A boundary state with these properties will be called irreducible. Two irreducible states differing only in the order of the angles ν_i are equivalent. Note that if $|\,\nu_1,\nu_2,...,\nu_n\rangle$ is an irreducible state then so is $|\,\nu_1,\nu_2,...,\nu_{k-1},\nu_{k+1},...,\nu_n\rangle$ obtained by removing ν_k.

Proposition 1 can be derived by induction on the level of boundary states using Proposition 2.

Proposition 2: If an incoming particle of rapidity $i\nu_{n+1}$ fuses with the irreducible state $|\,\nu_1,\nu_2,...,\nu_n\rangle$, then three cases can be distinguished:

1st case: $p(i\nu_{n+1})nFp(i\nu_k)$ and $p(i\nu_{n+1})nFp(-i\nu_k)$ for $k=1..n$. The produced state is equivalent to the level $n+1$ state $|\,\nu_1,...,\nu_n,\nu_{n+1}\rangle$, which is irreducible. ν_{n+1} has to satisfy (42) in this case in order for the reflection factors on the produced state to have no poles on the imaginary axis in the physical strip.

2nd case: $p(i\nu_{n+1})Fp(i\nu_k)$ for one or more values of k, $1\leq k\leq n$. Let ν_{k_0} be the greatest among the corresponding ν_k-s. The produced state is equivalent to $|\,\nu_1, ..., \nu_{k_0-1}, \nu'_{k_0}, \nu_{k_0+1}, ...\nu_n\rangle$, where ν'_{k_0} can be computed by fusing $p(i\nu_{n+1})$ and $p(i\nu_{k_0})$ (see fig. 9a.). It is easy to verify that ν'_{k_0} satisfies (42), so $|\,\nu_1, ..., \nu_{k_0-1}, \nu'_{k_0}, \nu_{k_0+1}, ...\nu_n\rangle$ has the property that $p(i\nu_i)nFp(i\nu_j)$ for $i, j = 1..n$ and $\det(R(i\nu'_{k_0})) \neq 0$, but $p(i\nu_i)nFp(-i\nu_j)$ is not necessarily true. If the latter property is also true for $|\,\nu_1, ..., \nu_{k_0-1}, \nu'_{k_0}, \nu_{k_0+1}, ...\nu_n\rangle$ then it is irreducible, otherwise the 3rd case applies to it with $p(i\nu'_{k_0})$ being the incoming bulk particle and $|\,\nu_1, ..., \nu_{k_0-1}, \nu_{k_0+1}, ...\nu_n, B_{\frac{1}{2}}\rangle$ the boundary state.

3rd case: $p(i\nu_i)nFp(i\nu_j)$ for $i, j = 1..n$, but $p(i\nu_{n+1})Fp(-i\nu_k)$ is true for certain values of k. Let ν_{k_0} be the smallest of the corresponding angles. The produced state is equivalent to $|\,\nu_1, ..., \nu_{k_0-1}, \nu'_{k_0}, \nu_{k_0+1}, ...\nu_n\rangle$, where ν'_{k_0} arises by fusing $p(i\nu_{n+1})$ and $p(-i\nu_{k_0})$ (see figs. 11 a.,b.). It is easy to verify that ν'_{k_0} satisfies (42), so this state has the property that $p(i\nu_i)nFp(i\nu_j)$ for $i, j = 1..n$. $\det(R(i\nu'_{k_0})) \neq 0$ is also true, but $p(i\nu_i)nFp(-i\nu_j)$ is not necessarily. If the latter condition is also satisfied, then $|\,\nu_1, ..., \nu_{k_0-1}, \nu'_{k_0}, \nu_{k_0+1}, ...\nu_n\rangle$ is irreducible, otherwise the present case applies to it with $p(i\nu'_{k_0})$ being the incoming bulk particle and $|\,\nu_1, ..., \nu_{k_0-1}, \nu_{k_0+1}, ...\nu_n\rangle$ the boundary state.

Even if applying the 2nd or the 3rd case does not result in an irreducible state, the level of the involved boundary state decreases by one, so an irreducible state is obtained in a finite number of steps.

The equivalences stated in Proposition 2 in the second and third cases can be derived as follows: as in the previous subsections, one uses a multi-particle reflection amplitude (see e.g. the r.h.s of fig. 8) to determine the transformation properties of the produced state. In the corresponding diagram the scattering vertex of $p(i\nu_{n+1})$ and $p(i\nu_{k_0})$ should be split to a fusion and a decay vertex first, and then the fusing and decaying points should be moved far away in the past and future (using the Yang-Baxter equations). This should be done in such a way that those reflections which are not moved (with angles $i\nu_k$, $k \neq n+1, k_0$) should eventually take place between the fusing and decaying points. If these points are moved sufficiently far away, then because of the choice of k_0 all two-particle scatterings before the fusing point and after the decaying point are bijective, so only the part of the diagram that lies between these points is relevant.

The statements above can be used to determine the multiplicities and transformation properties of a boundary bound state in concretely given models in a finite number of steps. $|\,\nu_1, \nu_2, ..., \nu_k\rangle$ being the boundary state, the irreducible state equivalent to $|\,\nu_1, \nu_2\rangle$ should be determined first, then using the result the state equivalent to $|\,\nu_1, \nu_2, \nu_3\rangle$ should be determined, and so on.

4.3 Boundary states generated by kinks and particles

The statements and arguments in this subsection are similar to those in 4.2.4.

Proposition 1: A general boundary bound state is equivalent to a state $|\,\nu_1, \nu_2, ..., \nu_n\rangle$ which is such that ν_i satisfies condition (42) if $p_i(i\nu_i)$ is a particle, and $p_i(i\nu_i)nFp_j(i\nu_j)$ and

$p_i(i\nu_i)nFp_j(-i\nu_j)$ are satisfied for any $i,j=1..n$, $i\neq j$. There is at most one kink among the p_i-s. A state with these properties will be called irreducible. The multiplicity of these states is 2^n and they are equivalent to $|\,p_1(i\nu_1),p_2(i\nu_2),...,p_n(i\nu_n),B_{\frac{1}{2}}\rangle$. The reflection amplitudes on these states have no poles on the imaginary axis in the physical strip. Two irreducible states differing only in the order of the corresponding $p_i(i\nu_i)$-s are equivalent. Removing any angle from an irreducible state results in an irreducible state.

Proposition 1 can be derived by induction on the level of boundary states using Proposition 2.

Proposition 2: If an incoming particle of rapidity $i\nu_{n+1}$ fuses with an irreducible boundary state, then the following four cases can be distinguished:

1st case: the boundary state is $|\,\nu_1,\nu_2,...,\nu_n\rangle$, where all the p_i-s ($i=1..n$) are particles and p_{n+1} is also a particle. This situation is described in section 4.2.

2nd case: the boundary state is $|\,\nu_1,\nu_2,...,\nu_n\rangle$, where all the p_i-s ($i=1..n$) are particles and p_{n+1} is a kink. If $p(i\nu_{n+1})nFp(-i\nu_k)$ is satisfied for $k=1..n$, then the produced state $|\,\nu_1,...,\nu_n,\nu_{n+1}\rangle$ is irreducible. If $p(i\nu_{n+1})Fp(-i\nu_k)$ for some values of k, then let ν_{k_0} be the smallest among the corresponding ν_k-s. The produced state is equivalent to the state $|\,\nu_1,...,\nu_{k_0-1},\nu_{k_0+1},...\nu_n,\nu'_{k_0}\rangle$, where ν'_{k_0} can be computed by fusing $p(i\nu_{n+1})$ and $p(i\nu_{k_0})$, and $p(i\nu'_{k_0})$ is a kink. $|\,\nu_1,...,\nu_{k_0-1},\nu_{k_0+1},...\nu_n,\nu'_{k_0}\rangle$ is either irreducible or the present case applies to it with $p(i\nu'_{k_0})$ being the incoming bulk kink.

3rd case: the boundary state is $|\,\nu_1,\nu_2,...,\mu_m,...,\nu_n\rangle$, p_m is a kink and p_{n+1} is a particle. This is very similar to the first case. It is possible that the produced state is equivalent to the irreducible state $|\,\nu_1,\nu_2,...,\mu_m,...,\nu_n,\nu_{n+1}\rangle$. In this case (42) must be satisfied for ν_{n+1}. The other possibility is that the produced state is equivalent to the state $|\,\nu_1,...,\nu_{k_0-1},\nu_{k_0+1},...\nu_n,\nu'_{k_0}\rangle$, where p_{k_0} is either a kink or a particle. This state is either irreducible or the present or the 2nd case applies to it with p_{k_0} being the incoming particle.

4th case: The boundary state is $|\,\nu_1,\nu_2,...,\mu_m,...,\nu_n\rangle$, where p_m is a kink and the incoming particle p_{n+1} is also a kink. In this case the vertex of the two kinks $p(i\mu_m)$ and $p(i\nu_{n+1})$ can be split and the fusing and decaying points should be moved far away to the past and future. Thus the produced state is equivalent to $|\,\nu_1,\nu_2,...,\nu'_m,...,\nu_n\rangle$, where $p(i\nu'_m)$ is a particle, ν'_m is determined by μ_m and ν_{n+1}. Now the first case can be applied with $p(i\nu'_m)$ being the incoming bulk particle and $|\,\nu_1,\nu_2,...,\nu_{m-1},\nu_{m+1},...,\nu_n\rangle$ being the boundary state. It is easy to see that ν' satisfies the condition (42) in a stronger form:

$$\nu'\leq u'. \tag{44}$$

The statements above can be used to determine the multiplicities and transformation properties of a boundary bound state in concretely given models in a finite number of steps (see also the end of section 4.2.4).

An important simple special case is the following: if $|\,\mu_1,\mu_2,...,\mu_n\rangle$ is such that all the p_i-s ($i=1..n$), are kinks and $\mu_i\neq\mu_j$ for all $i\neq j$, then the multiplicity of this state is the maximal $2^{\lceil n/2\rceil}$. This follows immediately from the bijectivity of S_K, but could also be derived using the propositions above. The reflection amplitudes on such a state have no poles on the imaginary axis in the physical strip.

We remark that irreducible states that are not equal in our terminology can be equivalent.

In the next subsection we bring the results of the present and 4.2 subsections to a simpler form.

4.4 Supersymmetric boundary states and fusing rules

To a supersymmetric particle (multiplet) $p(\theta)$ with mass $m = 2M\cos(\rho/2)$ we assign the following (not ordered) set of rapidities: $L_{p(\theta)} = \{\theta - i\rho/2, \theta + i\rho/2\}$, where it is not required that $\theta \pm i\rho/2$ be in the physical strip. The elements of the set are the rapidities of those kinks which fuse to the particle $p(\theta)$. $L_{p(\theta)}$ determines $p(\theta)$ uniquely. We assign the set $L_{k(\theta)} = \{\theta\}$ to a kink $k(\theta)$. This notation allows us to summarize the bulk supersymmetric fusing rules in a simple form. In terms of these sets the bulk fusion $p_1 + p_2 \to p_3$ of two particles takes the form

$$\{\theta_1, \theta_2\} + \{\theta_3, \theta_1 \pm i\pi\} \to \{\theta_2, \theta_3\}, \tag{45}$$

where $L_{p_1} = \{\theta_1, \theta_2\}$, $L_{p_2} = \{\theta_3, \theta_1 \pm i\pi\}$, $L_{p_3} = \{\theta_2, \theta_3\}$, and $\theta_1, \theta_2, \theta_3$ are appropriate complex rapidities. Similarly, the fusion of a kink and a particle $k_1 + p \to k_2$ takes the form

$$\{\theta_1\} + \{\theta_2, \theta_1 \pm i\pi\} \to \{\theta_2\}. \tag{46}$$

A kink-kink fusion $k(\theta_1) + k(\theta_2) \to p$ takes the form

$$\{\theta_1\} + \{\theta_2\} \to \{\theta_1, \theta_2\}. \tag{47}$$

The essential point is that in these fusions the set of rapidities corresponding to the final state is obtained in the following way: the disjoint union of the two sets of rapidities corresponding to the fusing particles/kinks is formed and the pair of rapidities differing by $\pm i\pi$ are deleted (if there is any such pair). We allow here and further on that a set contains certain elements several times, i.e. the elements of the sets we consider have multiplicity.

These rules are in accordance with the fact that S_K is bijective (of rank 6) in the physical strip but is of rank 3 at $\theta = i\pi$. The fusing rules (45), (46), (47) apply in the same form if we do not perform the factorizations described in section 2.3, i.e. if two kinks fuse into a state with multiplicity 6, which we call "two-kink-state". Correspondingly, the fusing rules given in section 2.3 also apply in this case with the minor modification that "two-kink-states" should be substituted for "supersymmetric particles". In this case the fusion of two kinks produces a "two-kink-state" state with multiplicity 6 as described in sections 2.2 and 2.3. It does not seem to be useful to stress the decomposition (8) because of the adjacency conditions and relations following from the kink adjacency conditions. Numerical (TCSA) calculations in finite volume suggest [20] that it is possible that the breathers of the bulk supersymmetric sine-Gordon model transform as “two-kink-states” rather than simple SUSY particles [17].

Let us assume that the particle reflections are described by the factors $R^{(\pm)}_{P1}$ or $R^{(-)}_{P2,e}$, i.e. by the factors that can be obtained by bootstrap from the kink reflection factors. The case of $R^{(+)}_{P2,e}$ will be considered later.

To a general boundary state $v = |\,\nu_1,\nu_2,...\nu_n\rangle$ we assign the set P_v the elements of which are the sets of rapidities assigned to $p_1, p_2, ..., p_n$ with the modification that all the rapidities $i\theta$ ($\theta \in \mathbb{R}$) are changed to $i|\theta|$. Note that $0 \leq |\theta| < \pi$. For example, $P_v = \{\{i|\theta_{11}|, i|\theta_{12}|\}, \{...\}, ...\}$ if p_1 is a particle and $L_{p_1(i\nu_1)} = \{i\theta_{11}, i\theta_{12}\}$. The disjoint union of the sets contained by P_v is denoted by $\cup P_v$. It is easily verified that v is irreducible if and only if $\cup P_v$ does not have two elements $i\theta_1, i\theta_2$ for which $\theta_1 + \theta_2 = \pi$.

Let v' denote the (not necessarily unique) irreducible state obtained from the boundary state v by applying the results of section 4.2 and 4.3. It is straightforward to verify that the effect of the steps (described in 4.2.4 and 4.3) used to obtain v' is that pairs of rapidities in $\cup P_v$ with the property $\theta_1 + \theta_2 = \pi$ are removed. Consequently, $\cup P_{v'}$ is obtained from $\cup P_v$ by removing all such pairs. We introduce the notation $L_v = \cup P_{v'}$.

Using BBE I it is easy to see that an irreducible state w is equivalent to the state $\tilde{w} = |\,\theta_1, \theta_2, ..., \theta_n\rangle$, where $p(i\theta_i)$ is a kink for all i-s, $L_w = \cup P_w = \{i\theta_i \,|\, i = 1...n\}$, and $0 < \theta_1 \leq \theta_2 \leq ... \leq \theta_n$. However, some of the θ-s can be outside the physical strip here. This state has multiplicity $2^{\lceil n/2 \rceil}$ and it is equivalent to $|\,k(i\theta_1), k(i\theta_2), ..., k(i\theta_n), B_{\frac{1}{2}}\rangle$.

Now it is clear that two irreducible states w_1 and w_2 are equivalent if $L_{w_1} = L_{w_2}$. This condition is also necessary: the bootstrap equations applied to $\tilde{w}$ (where w is an irreducible state) imply that the one-kink reflection amplitude $r(i\theta)$ on w can also be expressed as a product containing the factors $S_K(i(\theta \pm \theta_i)$, $R_K(i\theta)$ and θ-independent numerical factors, where $i\theta$ is the rapidity of the reflecting kink. Thus the sequence θ_i, $i = 1...n$ is uniquely determined by the analytic properties of $r(i\theta)$, so if w_1 and w_2 are equivalent, then $L_{w_1} = L_{w_2}$ must hold.

The supersymmetric boundary fusion rules can be summarized as follows: a supersymmetric boundary state v is equivalent to the state $\widetilde{(v')}$. Two supersymmetric boundary bound states v, z are equivalent if and only if $L_w = L_z$ (or equivalently $\widetilde{(v')} = \widetilde{(z')}$). The reflection factors on v have no poles and zeroes on the imaginary axis in the physical strip if and only if

$$\theta_i < \pi - \rho_{max}/2 \quad \forall i = 1...n. \tag{48}$$

(See (39) for ρ_{max}, $L_v = \{i\theta_i \,|\, i = 1...n\}$) Consider now the boundary fusion $p + v \to y$, where p is a bulk particle, v and y are boundary states. L_y can be obtained from L_p and L_v in the following way: the disjoint union $L_p \cup L_v$ should be formed, the rapidities $i\theta \in L_p \cup L_v$ should be replaced by the values $i|\theta|$, and all the pairs $i\theta_1, i\theta_2$ satisfying the condition $\theta_1 + \theta_2 = \pi$ should be removed. This rule is analogous to the bulk fusion rules, but the amplitudes $i\theta$ and $-i\theta$ ($\theta \in \mathbb{R}$) are identified in this case.

It is also possible that (48) is not satisfied for certain bound states, but the poles of the (supersymmetric) reflection factors on them are canceled by zeroes of the corresponding non-supersymmetric reflection factor, so the poles of the supersymmetric reflection factors do not introduce new bound states. The fusion rules above apply in this case as well.

These boundary fusion rules can be applied unchanged if instead of supersymmetric particles "two-kink-states" are taken.

The statements above also apply to the case when particle reflections on the ground state are described by the factor $R^{(+)}_{P2,e}$ with the following modification: the rapidities should not be altered when P_v is constructed, i.e. for example $P_v = \{\{i\theta_{11}, i\theta_{12}\}, \{...\}, ...\}$. Although a rapidity θ in $\cup P_v$ is not identified with $-\theta$, $\cup P_{v_1}$ and $\cup P_{v_2}$ are regarded equivalent if they can be obtained from one another by changing the sign of an even number of rapidities. L_v is obtained from $\cup P_v$ by removing all the pairs of amplitudes of the form $i\theta, -i\theta$, taking into consideration that the sign of an even number of amplitudes in $\cup P_v$ can be changed freely. If a change of signs gives rise to pairs of the form $i\theta, -i\theta$, then this change should be done, and the arising pairs should be removed. w is equivalent to $|\, k(i\theta_1), k(i\theta_2), ..., k(i\theta_n), B_{\frac{1}{2}}\rangle$, where $L_w = \{i\theta_1, ..., i\theta_n\}$. The states v and w are equivalent if and only if L_v can be transformed into L_w by changing the sign of an even number of rapidities. We use the notation $L_v \sim L_w$ in this case. L_y is obtained from $L_p \cup L_v$ in the same way as L_v from $\cup P_v$.

These modifications are due to the fact that $R^{(+)}_{P2,e}$ is not obtained from a kink amplitude by bootstrap and the boundary Yang-Baxter equation is not satisfied by $R^{(+)}_{P2,e}$ and $R^{(+)}_{K}$.

We consider now the case when kink rapidities $i(\pi - |\xi|)$ are also allowed, and the reflections on the ground state are described by the factors $R^{(-)}_{K,e}$, $R^{(-)}_{P2,e}$ or $R^{(+)}_{P2,e}$. This is the case when the ground state reflection factors are allowed to be degenerate at particular fusing rapidities (see also the first paragraph of section 4.2). The description above is modified in this case as follows: it is not allowed to change the sign of all the rapidities $i(\pi - |\xi|)$ in a set in the $(+)$ case.

In the $(-)$ case, if L_v contains $i(\pi - |\xi|)$, then the state v is equivalent to the state $|\, k(i\theta_1), k(i\theta_2), ..., k(i\theta_{l-1}), k(i\theta_{l+1}), ..., k(i\theta_n), B_{0/1}\rangle$, where $i\theta_1, ..., i\theta_n$ are the elements of L_v (they are not necessarily different), $\theta_l = \pi - |\xi|$, and $|\, B_{0/1}\rangle = |\, \pi - |\xi|\rangle$. It is allowed that $|\theta_j| = \pi - |\xi|$ for several integers $j \neq l$. The multiplicity of v is $2^{\lceil n/2\rceil - 1}$. $|\, B_{0/1}\rangle$ is a singlet state.

In the $(+)$ case v is equivalent to $|\, k(i\theta_1), k(i\theta_2), ..., k(i\theta_{l-1}), k(i\theta_{l+1}), ..., k(i\theta_{j-1}), k(i\theta_{j+1}), ..., k(i\theta_n), B'_{\frac{1}{2}}\rangle$, where j is chosen arbitrarily and $|\, B'_{\frac{1}{2}}\rangle = |\, \nu'\rangle$, $p(i\nu')$ is the particle constituted by the kinks $k(i\theta_l)$ and $k(i\theta_j)$. $|\, B'_{\frac{1}{2}}\rangle$ is a singlet state.

The rules and statements of the present subsection are sufficient to determine the transformation and scattering properties of the boundary bound states in concretely given models without the use of the results described in 4.2.4 and 4.3.

Further supersymmetric boundary theories may be obtained from known ones by projection, i.e. by factoring out certain boundary bound states. We do not discuss this possibility in detail.

It is important to note that we have not taken into consideration the "statistics" of the various bulk particles at all, i.e. we have assumed "free" statistics. The correct statistical properties can be achieved by projection. An example of this can be found in section 5.3.

Let us consider now the situation that a boundary state in the non-supersymmetric theory can be created in more than one way, i.e. the states v and v' are equivalent (and so regarded to be the same) in the non-supersymmetric theory. Such an equivalence is usually established by transforming certain diagrams representing amplitudes into each other, where the allowed transformations include shifting lines using bootstrap and Yang-Baxter equations and splitting or fusing vertices. These transformations are similar to those done above, and we expect that they can usually be used to establish the equivalence of the corresponding supersymmetric states as well. Therefore we expect that the supersymmetric states Sv and Sv' corresponding to v and v' are also equivalent, i.e. $L_{Sv} = L_{Sv'}$ (or $L_{Sv} \sim L_{Sv'}$). The equivalence of such states should be checked in specific models, but we expect that the requirement that such states should be equivalent is usually satisfied and does dot place constraints on the boundary fusing rules of the non-supersymmetric theory. The only really restrictive condition on the applicability of the Ansatz described in the previous sections is thus (48).

5 Examples

The following statement illustrates the phenomenon also encountered in the bulk scattering theory [3, 1] that prescribing the representations in which the boundary states transform places very restrictive conditions on the boundary fusing rules. Let P be a supersymmetric particle with parameters $m = 2M\sin(u)$ (where $M > 0$, $0 < u \leq \pi/2$). Let the boundary state $w = |\,\nu_1, \nu_2, ..., \nu_n\rangle$ be generated by this particle in n steps and assume that the ground state particle reflection factor is nondegenerate at $i\nu_i$, $i = 1..n$. In this case $|\,\nu_1\rangle$, $|\,\nu_1, \nu_2\rangle$,...,w have multiplicity 2 (and are equivalent to first level states) if and only if $\nu_k - \nu_{k-1} = 2u$ $\forall k = 2...n$.

For the sake of simplicity, we do not consider the case when particle reflections on the ground state are described by $R^{(+)}_{P2,e}$ in the next subsections.

5.1 Boundary sine-Gordon model

The particle spectrum of the bulk sine-Gordon theory (SG) contains a soliton (s) and an antisoliton ($\bar{s}$) of mass M and the breathers B_n of mass $m_n = 2M\sin(u_n)$, where

$u_n = \pi n/(2\lambda)$, $n = 1, ..., [\lambda]$, λ and M are parameters of the model. The breathers are self-conjugate, and the conjugate of s is $\bar{s}$. The fusing rules (given in the form process, (fusing angle)) are the following: $s + \bar{s} \to B_n$, $(\pi - 2u_n)$; $B_n + B_m \to B_{n+m}$, $(u_n + u_m)$ provided $n + m \leq [\lambda]$; and the crossed versions of these rules. They are consistent with associating kink representations to s and $\bar{s}$ and particle or "two-kink-state" representations with $\alpha = -1/(2M)$ to the breathers. The sets corresponding to the supersymmetric parts of s, $\bar{s}$, and B_n are $L_s = \{\theta\}$, $L_{\bar{s}} = \{\theta\}$, $L_{B_n} = \{\theta + i(\pi/2 - u_n), \theta - i(\pi/2 - u_n)\}$.

The boundary SG model (BSG) as defined in [19] (see also [18]) has the boundary spectrum containing the states $|\, n_1, n_2, ..., n_k\rangle$, where $n_1, n_2, ..., n_k$ are nonnegative integers satisfying the condition $\pi/2 \geq \nu_{n_1} > w_{n_2} > \nu_{n_3} > ... \geq 0$, where $\nu_n = \eta/\lambda - u_{2n+1}$, $w_n = \pi - \eta/\lambda - u_{2n-1}$, and $0 < \eta \leq \frac{\pi}{2}(\lambda + 1)$ is a boundary parameter. The fusing rules [18, 22] are listed in the following table:

Initial state	particle	rapidity	final state
$\|n_1, ..., n_{2k}\rangle$	$s, \bar{s}$	$i\nu_n$	$\|n_1, ..., n_{2k}, n\rangle$
$\|n_1, ..., n_{2k-1}\rangle$	$s, \bar{s}$	iw_n	$\|n_1, ..., n_{2k-1}, n\rangle$
$\|n_1, ..., n_{2k}, n_{2k+1}, ...\rangle$	B^n	$i\frac{1}{2}(\nu_l - w_{n-l})$	$\|n_1, ..., n_{2k}, l, n-l, n_{2k+1}, ...\rangle$
$\|n_1, ..., n_{2k-1}, n_{2k}, ...\rangle$	B^n	$i\frac{1}{2}(w_l - \nu_{n-l})$	$\|n_1, ..., n_{2k-1}, l, n-l, n_{2k}, ...\rangle$
$\|n_1, ..., n_{2k}, ...\rangle$	B^n	$i\frac{1}{2}(\nu_{-n_{2k}} - w_{n+n_{2k}})$	$\|n_1, ..., n_{2k} + n, ...\rangle$
$\|n_1, ..., n_{2k-1}, ...\rangle$	B^n	$i\frac{1}{2}(w_{-n_{2k-1}} - \nu_{n+n_{2k-1}})$	$\|n_1, ..., n_{2k-1} + n, ...\rangle$

The first two lines show that the whole boundary spectrum can be generated by kinks. Correspondingly, we associate to the BSG state $|\, n_1, n_2, ..., n_k\rangle$ the supersymmetric part $|\, \nu_{n_1}, w_{n_2}, \nu_{n_3}, ... B_{\frac{1}{2}}\rangle$ (using the notation introduced earlier), where $p(i\nu_{n_1})$, $p(iw_{n_2})$... are kinks. Now we have to verify if the fusion rules given in the 3-6th lines are also valid for these supersymmetric parts. This is easily done by using the rules given in section 4.4. Let us consider the 3rd line first. Let $v = |\, \nu_{n_1}, ..., w_{n_{2k}}, \nu_{n_{2k+1}}, ..., B_{\frac{1}{2}}\rangle$, $p = p(i\frac{1}{2}(\nu_l - w_{n-l}))$ with u_n, $w_{2k} > \nu_l > w_{n-l} > \nu_{2k+1}$, and $p + v \to y$. In this case $L_p = \{i\nu_l, -iw_{n-l}\}$, so $L_y = \{i\nu_{n_1}, \ldots, iw_{n_{2k}}, i\nu_{n_{2k+1}}, \ldots, i\nu_l, iw_{n-l}\}$. The 4th line is similar. Turning to the 5th line, let $v = |\, \nu_{n_1}, ..., w_{n_{2k}}, ..., B_{\frac{1}{2}}\rangle$, $p = p(i\frac{1}{2}(\nu_{-n_{2k}} - w_{n+n_{2k}}))$ with u_n, and $p + v \to y$. Now $L_p = \{i\nu_{-n_{2k}}, -iw_{n+n_{2k}}\}$, and because of $\nu_{-n_{2k}} + w_{n_{2k}} = \pi$ we have $L_y = \{i\nu_{n_1}, ..., iw_{n+n_{2k}}, ...\}$. The 6th line is similar to the 5th line.

Condition (48) is clearly satisfied for all boundary states, so the supersymmetric factors of the reflection amplitudes have no poles on the imaginary axis in the physical strip.

We remark that the one-particle reflection amplitudes and two-particle S-matrices of the BSG model contain two bulk and two boundary parameters. The relation between these parameters and the parameter γ (see (33)) in the $(-)$ case is not yet known, so it cannot be decided whether any of the angles ν_n and w_n coincides with $\pi - \xi$ or not. The transformation and scattering properties of the boundary states are modified by such a coincidence.

Because of the solitons the factor $R_{P2,e}^{(+)}$ cannot be used to describe the reflections of breathers on the ground state boundary. This factor can be used, however, if one restricts to the breather sector of the model.

5.2 Boundary sinh-Gordon model

The boundary sinh-Gordon (BShG) model is obtained from the BSG model by analytic continuation in the bulk coupling constant β, where β is related to λ through $\lambda = \frac{8\pi-\beta^2}{\beta^2}$. In ShG $\beta = i\hat{\beta}$, where $\hat{\beta} \in \mathbb{R}$, so $\lambda = -\frac{8\pi+\hat{\beta}^2}{\hat{\beta}^2} < -1$. The particle spectrum of the ShG model contains only one self-conjugate particle P with a two-particle scattering amplitude and reflection amplitude that can also be obtained from the corresponding amplitudes of the first SG breather (B_1) by the analytic continuation $\beta = i\hat{\beta}$. BShG also contains a series of boundary states b_n [12] corresponding to the BSG states $|\,\nu_0, w_n\rangle$, $n = 1...$ which are precisely those states that can be generated using B_1 only. b_0 is the ground state. The bulk fusing rules are the following:

Initial state	particle	rapidity	final state
b_n	P	$i(\frac{\eta}{\lambda} - \frac{\pi}{2} + \frac{\pi}{\lambda}n)$	b_{n+1}

The supersymmetric BShG model is also obtained by the analytic continuation above. This implies that in the formula $m = 2M\sin(u)$ for the particle mass $M < 0$ and $u < 0$, i.e. the values of these parameters are not in the range that is considered in our paper. A consequence of this, for example, is that the supersymmetric factor of the two-particle scattering amplitude has a pole in the physical strip (canceled by a zero of the non-supersymmetric factor) [11]. However, the supersymmetric parts of the states b_n and the corresponding reflection factors can be obtained by the same steps as those of $|\,\nu_0, w_n\rangle$ but with $M < 0$, $u < 0$, so the diagrams cannot be drawn as certain angles take non-physical values. Thus the states b_n have multiplicity two in the supersymmetric BShG model if the value of ξ is generic. The situation is similar if the ground state reflection factor is $R_{P2,e}^{(+)}$.

5.3 Free particle on the half line

The supersymmetric factors of the scattering and ground state reflection amplitudes of this model can be obtained by taking the limit $\alpha \to 0$, which implies $M \to \infty$, $u \to 0$, $\rho \to \pi$, in the $(-)$ case $\xi \to \pi$, $R_{P2,e}^{(-)} \to R_{P1}^{(-)}$, and in the $(+)$ case $\xi \to 0$, $R_{P2,e}^{(+)} \to R_{P1}^{(+)}$. The same result can be obtained by solving the Yang-Baxter equations. S_P is also bijective in the physical strip. The arguments used to establish the results described in section 4.2.4 can be applied to this case as well, and it is easy to see that the multiplicity of a level n boundary bound state is 2^n irrespectively of the particular boundary fusing rules. The reflection amplitudes on the boundary bound states are also free from physical strip poles for any values of the fusing angles. However, if some value of the boundary fusing angles occurs several times, then the statistical properties of the particles should be taken into consideration. In particular, if all the boundary fusing angles are the same (see [12]), then the multiplicity of a level n boundary bound state is 2 for each value of n. This is consistent with the multiplicities obtained by taking the zero bulk coupling limit of the boundary sinh-Gordon model.

5.4 Boundary $a_2^{(1)}$ affine Toda field theory

The bulk spectrum of this model contains two particles 1 and 2 of equal mass $m_1 = m_2$. Their fusing rules are $1+1 \to 2$ $(\pi/3)$ and $2+2 \to 1$ $(\pi/3)$. The charge conjugate of 1 is 2. These rules are easily seen to be consistent with associating supersymmetric particle representations to 1 and 2 with $\alpha = -1/(2M)$, $m_1 = m_2 = 2M\sin(\pi/3)$ [1]. M is a parameter of the model.

The boundary version of this model described in [13] contains the boundary states $b_{n,m}$ for all $n, m \in \mathbb{Z}$, $n+m \geq 0$, $-\frac{1}{2B} - \frac{1}{2} < n, m < \frac{1}{2B} + \frac{1}{2}$, and the states $b_{-n,n}$ and $b_{n,-n}$ for all $n \in \mathbb{Z}$, $0 \leq \frac{3}{2B} + \frac{1}{2}$, where B is a parameter of the bulk model. We consider only generic values of B and only the domain $0 < B < 1$, as in [13]. $b_{0,0}$ is the ground state. The fusing rules are shown in the following table:

Initial state	particle	rapidity	final state
$b_{n,m}$	1	$i\frac{\pi}{6} - i\frac{\pi}{6}B(2n+1)$	$b_{n+1,m}$
$b_{n,m}$	2	$i\frac{\pi}{6} - i\frac{\pi}{6}B(2m+1)$	$b_{n,m+1}$
$b_{-n,n}$, $0 \leq n$	1	$i\frac{\pi}{2} - i\frac{\pi}{6}B(2n+1)$	$b_{-n-1,n+1}$
$b_{n,-n}$, $0 \leq n$	2	$i\frac{\pi}{2} - i\frac{\pi}{6}B(2n+1)$	$b_{n+1,-n-1}$

The supersymmetric parts associated to these boundary bound states are denoted by $Sb_{n,m}$, $Sb_{-n,n}$, $Sb_{n,-n}$ respectively. It is straightforward to verify that the fusing rules above imply that the corresponding sets $L_{Sb_{n,m}}$, $L_{Sb_{-n,n}}$, $L_{Sb_{n,-n}}$ are the following:

$L_{Sb_{n,m}} = \{i(\frac{\pi}{3} - \frac{\pi}{6}B(2j+1)), i(\frac{\pi}{6}B(2j+1)), i(\frac{\pi}{3} - \frac{\pi}{6}B(2l+1)), i(\frac{\pi}{6}B(2l+1)) \,|\, j = 1...n, l = 1...m\}$ if $n, m \geq 0$,

$L_{Sb_{-n,n}} = L_{Sb_{n,-n}} = \{i(\frac{2\pi}{3} - \frac{\pi}{6}B(2j+1)), i(\frac{\pi}{3} - \frac{\pi}{6}B(2j+1)) \,|\, j = 1...n\}$,

$L_{Sb_{n,m}} = \{i(\frac{2\pi}{3} - \frac{\pi}{6}B(2j+1)), i(\frac{\pi}{3} - \frac{\pi}{6}B(2l+1)), i(\frac{\pi}{6}B(2r+1)) \,|\, j = 1...(-n), l = 1...m, r = (-n+1)...m\}$ if $n < 0$,

$L_{Sb_{n,m}} = \{i(\frac{2\pi}{3} - \frac{\pi}{6}B(2j+1)), i(\frac{\pi}{3} - \frac{\pi}{6}B(2l+1)), i(\frac{\pi}{6}B(2r+1)) \,|\, j = 1...(-m), l = 1...n, r = (-m+1)...n\}$ if $m < 0$.

Consequently, the multiplicity of $Sb_{n,m}$ is 2^{n+m} if $n, m \geq 0$, 2^m if $n < 0$, 2^n if $m < 0$, provided that there is no coincidence between $\pi - \xi$ and the rapidities above. $\pi - \rho_{max}/2 = 5\pi/6$, so condition (48) is satisfied and the supersymmetric factors of the reflection amplitudes have no poles on the imaginary axis in the physical strip.

5.5 Boundary $a_4^{(1)}$ affine Toda field theory

The bulk spectrum of this model contains four particles 1, 2, 3, 4 of mass $m_1 = m_4 = 2M\sin(\pi/5)$, $m_2 = m_3 = 2M\sin(2\pi/5)$, where M is a parameter of the model. The fusing rules are $a+b \rightarrow c$, where either $c = a+b$ or $c = a+b-5$. The corresponding fusing angles are $\frac{\pi}{5}(a+b)$ if $c = a+b$ and $\frac{\pi}{5}(10-a-b)$ if $c = a+b-5$. The charge conjugate of 1 and 2 is 4 and 3. Theses rules are easily seen to be consistent with associating supersymmetric particle representations to 1, 2, 3, 4 with $\alpha = -1/(2M)$ [1].

The boundary version of this model described in [13] has two classes of inequivalent solitonic boundary conditions to which different boundary spectra belong.

If the boundary conditions are of the first class, then there are the boundary states b_{n_2,n_3}, $n_2, n_3 \in \mathbb{Z}$, $n_2 + n_3 \geq 0$, $-\frac{1}{2B} - \frac{1}{2} < n_2, n_3 < \frac{1}{2B} + \frac{1}{2}$, and $b_{n,-n}$ and $b_{-n,n}$ for all $n \in \mathbb{Z}$, $0 \leq n < \frac{5}{2B} + \frac{1}{2}$, where B is a parameter of the bulk model. Generic values of B are considered in the domain $0 < B < 1$. The fusing rules, that are analogous to those of the $a_2^{(1)}$ model are listed in the following table:

Initial state	particle	rapidity	final state
b_{n_2,n_3}	2	$i\frac{\pi}{10} - i\frac{\pi}{10}B(2n_2+1)$	b_{n_2+1,n_3}
b_{n_2,n_3}	3	$i\frac{\pi}{10} - i\frac{\pi}{10}B(2n_3+1)$	b_{n_2,n_3+1}
$b_{-n,n}$, $0 \leq n$	1	$i\frac{\pi}{2} - i\frac{\pi}{10}B(2n+1)$	$b_{-n-1,n+1}$
$b_{n,-n}$, $0 \leq n$	4	$i\frac{\pi}{2} - i\frac{\pi}{10}B(2n+1)$	$b_{n+1,-n-1}$

The supersymmetric parts associated to these particles are denoted by Sb_{n_2,n_3}, $Sb_{-n,n}$, $Sb_{n,-n}$ respectively. We have

$L_{Sb_{n_2,n_3}} = \{i(\frac{\pi}{5} - \frac{\pi}{10}B(2j+1)), i(\frac{\pi}{10}B(2j+1)), i(\frac{\pi}{5} - \frac{\pi}{10}B(2l+1)), i(\frac{\pi}{10}B(2l+1)) \,|\, j = 1...n_2, l = 1...n_3\}$ if $n_2, n_3 \geq 0$,

$L_{Sb_{-n,n}} = L_{Sb_{n,-n}} = \{i(\frac{4\pi}{5} - \frac{\pi}{10}B(2j+1)), i(\frac{\pi}{5} - \frac{\pi}{10}B(2j+1)) \,|\, j = 1...n\}$,

$L_{Sb_{n_2,n_3}} = \{i(\frac{4\pi}{5} - \frac{\pi}{10}B(2j+1)), i(\frac{\pi}{5} - \frac{\pi}{10}B(2l+1)), i(\frac{\pi}{10}B(2r+1)) \,|\, j = 1...(-n_2), l = 1...n_3, r = (-n_2+1)...n_3\}$ if $n_2 < 0$,

$L_{Sb_{n_2,n_3}} = \{i(\frac{4\pi}{5} - \frac{\pi}{10}B(2j+1)), i(\frac{\pi}{5} - \frac{\pi}{10}B(2l+1)), i(\frac{\pi}{10}B(2r+1)) \,|\, j = 1...(-n_3), l = 1...n_2, r = (-n_3+1)...n_2\}$ if $n_3 < 0$.

The multiplicity of Sb_{n_2,n_3} is $2^{n_2+n_3}$ if $n_2, n_3 \geq 0$, 2^{n_3} if $n_2 < 0$, 2^{n_2} if $n_3 < 0$, provided that there is no coincidence between $i(\pi - \xi)$ and the rapidities above.

If B is sufficiently small, then the states $Sb_{-1,1}, Sb_{-2,2}, ...$ violate condition (48), so the supersymmetric factor of the reflection amplitudes of particle 1 on these states have poles on the imaginary axis in the physical strip. There is one such pole for $Sb_{-1,1}$ which is canceled by a zero of the non-supersymmetric factor of the reflection amplitude, but there are two such poles for $Sb_{-2,2}$, of which only one is canceled. Applying lemma 1 in [21] it is not hard to see that the remaining pole cannot be explained by a Coleman-Thun mechanism. The SUSY factor of the reflection amplitude of 1 on $Sb_{-n,n}$ has in general n poles located at $i(\frac{\pi}{10} - \frac{\pi}{10}B(2m+1))$, $m = 1...n$ of which only the one at $i(\frac{\pi}{10} - \frac{3\pi}{10}B)$ is canceled by the non-supersymmetric factor, and they cannot be explained by Coleman-Thun mechanism. The situation is similar if the ground state reflection factor is $R^{(+)}_{P2,e}$.

If the boundary conditions are of the second class, then there are the boundary states b_{n_1,n_2,n_3,n_4}, $n_1, n_2, n_3, n_4 \in \mathbb{Z}$, $n_1 + n_2 \geq 0$, $n_2 + n_3 \geq 0$, $n_3 + n_4 \geq 0$. The fusing rules are the following:

Initial state	particle	rapidity	final state
b_{n_1,n_2,n_3,n_4}	1	$i\frac{\pi}{10} - i\frac{\pi}{10}B(2n_1+1)$	b_{n_1+1,n_2,n_3,n_4}
b_{n_1,n_2,n_3,n_4}	2	$i\frac{\pi}{10} - i\frac{\pi}{10}B(2n_2+1)$	b_{n_1,n_2+1,n_3,n_4}
b_{n_1,n_2,n_3,n_4}	3	$i\frac{\pi}{10} - i\frac{\pi}{10}B(2n_3+1)$	b_{n_1,n_2,n_3+1,n_4}
b_{n_1,n_2,n_3,n_4}	4	$i\frac{\pi}{10} - i\frac{\pi}{10}B(2n_4+1)$	b_{n_1,n_2,n_3,n_4+1}
b_{-n_2,n_2,n_3,n_4}, $0 \leq n_2$	1	$i\frac{3\pi}{10} - i\frac{\pi}{10}B(2n_2+1)$	$b_{-n_2-1,n_2+1,n_3 n_4}$
$b_{n_1,n_2,n_3,-n_3}$, $0 \leq n_3$	4	$i\frac{3\pi}{10} - i\frac{\pi}{10}B(2n_3+1)$	b_{n_1,n_2,n_3+1,n_3-1}

Using our rules we get

$L_{Sb_{n_1,n_2,n_3,n_4}} = \{i(\frac{2\pi}{5} - \frac{\pi}{10}B(2l_1+1)), i(\frac{\pi}{5} + \frac{\pi}{10}B(2l_1+1)), i(\frac{2\pi}{5} - \frac{\pi}{10}B(2l_4+1)), i(\frac{\pi}{5} + \frac{\pi}{10}B(2l_4+1)),$

$i(\frac{\pi}{5} - \frac{\pi}{10}B(2l_2+1)), i(\frac{\pi}{10}B(2l_2+1)), i(\frac{\pi}{5} - \frac{\pi}{10}B(2l_3+1)), i(\frac{\pi}{10}B(2l_3+1)) \,|\, l_i = 1...n_i\}$ if $n_1, n_2, n_3, n_4 \geq 0$,

$L_{Sb_{-n_2,n_2,n_3,n_4}} = \{i(\frac{\pi}{5} - \frac{\pi}{10}B(2l_3+1)), i(\frac{\pi}{10}B(2l_3+1)), i(\frac{2\pi}{5} - \frac{\pi}{10}B(2l_4+1)), i(\frac{\pi}{5} + \frac{\pi}{10}B(2l_4+1)),$

$i(\frac{6\pi}{10} - \frac{\pi}{10}B(2l_2+1)), i(\frac{\pi}{10}B(2l_2+1)) \,|\, l_3 = 1...n_3, l_4 = 1...n_4, l_2 = 1...n_2\}$ if $n_3, n_4 \geq 0$,

$L_{Sb_{n_1,n_2,n_3,-n_3}}$ is similar to $L_{Sb_{-n_2,n_2,n_3,n_4}}$,

$L_{Sb_{-n_2,n_2,n_3,-n_3}} = \{i(\frac{6\pi}{10} - \frac{\pi}{10}B(2l_2+1)), i(\frac{\pi}{10}B(2l_2+1)), i(\frac{6\pi}{10} - \frac{\pi}{10}B(2l_3+1)), i(\frac{\pi}{10}B(2l_3+1)) \,|\, l_2 = 1...n_2, l_3 = 1...n_3\}$,

$L_{Sb_{n_1,n_2,n_3,n_4}} = \{i(\frac{\pi}{5} - \frac{\pi}{10}B(2l_3+1)), i(\frac{\pi}{10}B(2l_3+1)), i(\frac{2\pi}{5} - \frac{\pi}{10}B(2l_4+1)), i(\frac{\pi}{5} + \frac{\pi}{10}B(2l_4+1)),$

$i(\frac{6\pi}{10} - \frac{\pi}{10}B(2j_1+1)), i(\frac{\pi}{10}B(2j_2+1)), i(\frac{\pi}{5} + \frac{\pi}{10}B(2j_3+1)) \,|\, l_3 = 1...n_3, l_4 = 1...n_4, j_2 = 1...n_2, j_1 = 1...(-n_1), j_3 = (-n_1+1)...n_2\}$ if $n_3, n_4, n_2 \geq 0$, $n_1 < 0$,

the remaining two cases when $n_4 < 0$, $n_1 \geq 0$ and $n_4, n_1 < 0$ are similar.

The multiplicity of Sb_{n_1,n_2,n_3,n_4} is $2^{n_1+n_2+n_3+n_4}$ if $n_1, n_2, n_3, n_4 \geq 0$, $2^{n_2+n_3+n_4}$ if $n_1 < 0$, $n_4 \geq 0$, $2^{n_1+n_2+n_3}$ if $n_1 \geq 0$, $n_4 < 0$ and $2^{n_2+n_3}$ if $n_1, n_4 < 0$ provided that there is no coincidence between $\pi - \xi$ and the rapidities above. Condition (48) is satisfied in this case, so the supersymmetric factors of the reflection amplitudes have no poles on the imaginary axis in the physical strip.

6 Discussion

We considered the boundary supersymmetric bootstrap programme in the case when the ground state is a singlet with RSOS label $\frac{1}{2}$ and the bulk particles transform in the kink and boson/fermion representations.

We presented a review of the relevant analogous results for bulk bootstrap. We introduced the boundary supersymmetry algebra and its action in the framework proposed by [15, 16], which requires that the boundary supersymmetry algebra has to be a coideal of the bulk SUSY algebra. In accordance with the literature, we found that there are two possible boundary supersymmetry algebras. The corresponding two cases —denoted by $(+)$ and $(-)$— lead to different supersymmetric ground state reflection factors. We found that these factors are essentially the same as those given in [14, 10, 9]. Although the algebras of the $(+)$ and $(-)$ cases appear to play symmetric role, the corresponding kink reflection factors turn out to be significantly different. A further important difference between the two cases is that in the $(+)$ case the boson/fermion reflection factor can be obtained by bootstrap from the kink reflection factor only at special values of its parameters [10]. We also found that the kink and boson/fermion reflection factors can be degenerate at particular rapidities depending on a parameter γ of the ground state representation.

We presented supersymmetric boundary fusing rules by which the representations and reflection factors for excited boundary bound states can be easily determined in specific models. The main difficulty of the problem is to handle the degeneracies of the boundary fusing tensors that occur at particular rapidities (resulting from the degeneracies of the one-particle reflection factors). These degeneracies are closely related to the degeneracies of the bulk two-particle scattering factors and of the ground state one-particle reflection factors. We found that the boundary fusing rules are analogous to the bulk rules [1, 3], and that it is useful to characterize the boson/fermion multiplets by their constituent kinks.

We considered the simplest case, when the two-particle scattering factors and ground state reflection factors are minimal and have no poles and overall zeroes on the imaginary axis in the physical strip, and there is no interplay between the poles and zeroes of the supersymmetric and non-supersymmetric factors of the S-matrix and reflection amplitude. We found that the main restriction on the applicability of the described construction follows from this condition and from the condition that the ground state is a singlet with

RSOS label $\frac{1}{2}$ (whereas in the bulk the main restriction arises because the supersymmetry representations are prescribed).

We applied our results to the boundary sine-Gordon model [18, 2], to the $a_2^{(1)}$ and $a_4^{(1)}$ affine Toda field theories [13] and to the free particle. We found that the boundary $a_2^{(1)}$ affine Toda model admits a tensor product type supersymmetrization, whereas the minimal supersymmetrization of the $a_4^{(1)}$ model with first class boundary condition is not possible: the supersymmetric reflection factors on some excited boundary states introduce poles that cannot be explained by Coleman-Thun mechanism. It is an open problem whether the supersymmetrized $a_2^{(1)}$ and $a_4^{(1)}$ reflection amplitudes describe any Lagrangian field theory. We also considered the sinh-Gordon model briefly.

It is likely that representations beyond the kink and boson/fermion for the bulk particles and other (possibly non-singlet) ground state representations can also be relevant to some models.

Acknowledgments

The author would like to thank L. Palla, G. Takács and Z. Bajnok for proposing this problem, for very useful discussions, and for comments on the manuscript.

Appendix

$$G^{[i,j]}(\theta)=R^{[i,j]}(\theta)R^{[i,j]}(\pi i-\theta)$$

$$R^{[i,j]}(\theta)=\frac{1}{\Gamma(\frac{\theta}{2\pi i})\Gamma(\frac{\theta}{2\pi i}+\frac{1}{2})}\prod_{k=1}^{\infty}\frac{\Gamma(\frac{\theta}{2\pi i}+\Delta_1+k+1)\Gamma(\frac{\theta}{2\pi i}-\Delta_1+k)}{\Gamma(\frac{\theta}{2\pi i}+\Delta_1+k-\frac{1}{2})\Gamma(\frac{\theta}{2\pi i}-\Delta_1+k+\frac{1}{2})}\times$$
$$\times\frac{\Gamma(\frac{\theta}{2\pi i}+\Delta_2+k-\frac{1}{2})\Gamma(\frac{\theta}{2\pi i}-\Delta_2+k-\frac{1}{2})}{\Gamma(\frac{\theta}{2\pi i}+\Delta_2+k)\Gamma(\frac{\theta}{2\pi i}-\Delta_2+k)},$$

where

$$\Delta_1=(u_i+u_j)/(2\pi),\qquad \Delta_2=(u_i-u_j)/(2\pi).$$

$$ZX^{(+)}(\theta)=\sqrt{m}P(\theta+i\rho/2)P(\theta-i\rho/2)\sqrt{2}K(2\theta)2^{-\theta/(i\pi)},$$

where $m=2M\cos(\frac{\rho}{2})$, $0\leq\rho<\pi$, $M=-1/\alpha$.

$$\tilde{Z}^{(-)}(\theta)=K(2\theta)2^{-\theta/(i\pi)}F(\theta-i\rho/2)F(\theta+i\rho/2),\qquad F(\theta)=P(\theta)K(\theta+i\xi)K(\theta-i\xi),$$

where $\gamma=-2\sqrt{M}\cos\frac{\xi}{2}$, $0\leq\xi\leq\pi$ or $Re(\xi)=0$ or $Re(\xi)=\pi$.

$$ZX^{(-)}(\theta)=\sqrt{m}\tilde{Z}^{(-)}(\theta)\frac{1}{\sqrt{2}}(\cos(\rho/2)-\cosh(\theta)),$$

where $\xi = \pi$.

$\tilde{Z}^{(+)}(\theta) = iK(2\theta)2^{-\theta/(i\pi)}F(\theta - i\rho/2)F(\theta + i\rho/2)U(\theta), \qquad F(\theta) = P(\theta)K(\theta + i\xi)K(\theta - i\xi),$

where $\gamma = -2\sqrt{M}i\sin(\xi/2)$, $-\pi \leq \xi \leq \pi$ or $Re(\xi) = 0$ or $Re(\xi) = \pm\pi$,

$$U(\theta) = f(\theta)/f(-\theta),$$

$$\frac{U(i\theta)}{U(i\theta - i\pi)} = -\frac{\cos(\theta) - \cos(\rho/2)}{\cos(\theta) + \cos(\rho/2)},$$

$$f(\theta) = \prod_{k=1}^{\infty} \frac{\Gamma(\frac{\rho}{4\pi} - \frac{\theta}{2\pi i} - \frac{1}{2} + k)\Gamma(\frac{\rho}{4\pi} + \frac{\theta}{2\pi i} + k)\Gamma(-\frac{\rho}{4\pi} - \frac{\theta}{2\pi i} - \frac{1}{2} + k)\Gamma(-\frac{\rho}{4\pi} + \frac{\theta}{2\pi i} + k)}{\Gamma(\frac{\rho}{4\pi} - \frac{\theta}{2\pi i} + k)\Gamma(\frac{\rho}{4\pi} + \frac{\theta}{2\pi i} + \frac{1}{2} + k)\Gamma(-\frac{\rho}{4\pi} - \frac{\theta}{2\pi i} + k)\Gamma(-\frac{\rho}{4\pi} + \frac{\theta}{2\pi i} + \frac{1}{2} + k)}.$$

$$K(\theta) = \frac{1}{\sqrt{\pi}} \prod_{k=1}^{\infty} \frac{\Gamma(k - \frac{1}{2} + \frac{\theta}{2\pi i})\Gamma(k - \frac{\theta}{2\pi i})}{\Gamma(k + \frac{1}{2} - \frac{\theta}{2\pi i})\Gamma(k + \frac{\theta}{2\pi i})},$$

$$P(\theta) = \prod_{k=1}^{\infty} \left[\frac{\Gamma(k - \frac{\theta}{2\pi i})^2}{\Gamma(k - \frac{1}{4} - \frac{\theta}{2\pi i})\Gamma(k + \frac{1}{4} - \frac{\theta}{2\pi i})} / \{\theta \leftrightarrow -\theta\} \right].$$

References

[1] T. J. Hollowood, E. Mavrikis: The N=1 supersymmetric bootstrap and Lie algebras, *Nucl. Phys.* B484 (1997) 631-652, hep-th/9606116.

[2] Z. Bajnok, L. Palla, G. Takács: Spectrum of boundary states in N=1 SUSY sine-Gordon theory, *Nucl. Phys.* B644 (2002) 509 hep-th/0207099

[3] K. Schoutens: Supersymmetry and factorized scattering, *Nucl. Phys.* B344 (1990) 665-695.

[4] K. Schoutens, M. Moriconi: Reflection matrices for integrable N=1 supersymmetric theories, *Nucl. Phys.* B487 (1997) 756-778, hep-th/9605219

[5] D. Bernard, A. LeClair: Residual quantum symmetries of the restricted sine-Gordon theories, *Phys. Lett.* B227:417, (1989)

[6] C. Ahn, D. Bernard, A. LeClair: Fractional supersymmetries in perturbed coset CFTs and integrable soliton theory, *Nucl. Phys.* B346 (1990) 409-439

[7] R. I. Nepomechie: The boundary supersymmetric sine-Gordon model revisited, *Phys. Lett.* B509 (2001) 183-188, hep-th/0103029

[8] R.I. Nepomechie: Supersymmetry in the boundary tricritical Ising field theory, *Int. J. Mod. Phys.* A17 3809 (2002), hep-th/0203123

[9] C. Ahn, W. Koo: Exact boundary scattering matrices of the supersymmetric sine-Gordon theory on a half line, *J. Phys.* A29 5845-5854, (1996), hep-th/9509056

[10] C. Ahn, W. Koo: Supersymmetric sine-Gordon model and the eight-vertex free fermion model with boundary, *Nucl. Phys.* B482 (1996) 675-695, hep-th/9606003

[11] C. Ahn, R. I. Nepomechie: Exact solution of the supersymmetric sinh-Gordon theory on a half line, *Nucl. Phys.* B586 (2000) 611-640, hep-th/0005170

[12] E. Corrigan, A. Taormina: Reflection factors and a two-parameter family of boundary bound states in the sinh-Gordon model, *J. Phys.* A33:8739-8754, 2000, hep-th/0008237

[13] G. W. Delius, G. M. Gandenberger: Particle reflection amplitudes in $a_n^{(1)}$ Toda field theories, *Nucl. Phys.* B554 (1999) 325-364, hep-th/9904002

[14] L. Chim: Boundary S-matrix for the tricritical Ising model, *Int. J. Mod. Phys.* A11 4491-4512, (1996), hep-th/9510008

[15] G. W. Delius, A. George: Quantum group symmetry of integrable models on the half-line, hep-th/0212300

[16] G. W. Delius, N. J. MacKay: Quantum group symmetry in sine-Gordon and affine Toda field theories on the half-line, hep-th/0112023

[17] C. Ahn: Complete S-matrices of supersymmetric sine-Gordon theory and perturbed superconformal minimal model, *Nucl. Phys.* B354 (1991) 57-84

[18] Z. Bajnok, L. Palla, G. Takács, G. Zs. Tóth: The spectrum of boundary states in sine-Gordon model with integrable boundary conditions, *Nucl. Phys.* B622 (2002) 548-564, hep-th/0106070

[19] S. Ghoshal, A. Zamolodchikov: Boundary S matrix and boundary state in two-dimensional integrable quantum field theory, *Int. J. Mod. Phys.* A9 (1994) 3841, hep-th/9306002

[20] Z. Bajnok, C. Dunning, L. Palla, G. Takács, F. Wágner: hep-th/0309120

[21] P. Mattsson, P. Dorey: Boundary spectrum in the sine-Gordon model with Dirichlet boundary conditions, *J. Phys.* A33:9065-9094, (2000), hep-th/0008071

[22] P. Mattsson: Integrable quantum field theories in the bulk and with a boundary, PhD thesis, University of Durham (2000), hep-th/0111261